arXiv:1405.1784v1 [math.AP] 8 May 2014

TIME DECAY OF SCALING INVARIANT ELECTROMAGNETIC SCHRÖDINGER EQUATIONS ON THE PLANE

LUCA FANELLI, VERONICA FELLI, MARCO A. FONTELOS, AND ANA PRIMO

ABSTRACT. We prove the sharp $L^1 - L^\infty$ time-decay estimate for the $2D$ - Schrödinger equation with a general family of scaling critical electromagnetic potentials.

1. INTRODUCTION

Let us consider an electromagnetic Schrödinger equation of the type

$$iu_t = \left(-i\nabla + \frac{\mathbf{A}\left(\frac{x}{|x|}\right)}{|x|}\right)^2 u + \frac{a\left(\frac{x}{|x|}\right)}{|x|^2}\, u, \tag{1.1}$$

where $N \geqslant 2$, $u = u(x,t) : \mathbb{R}^{N+1} \to \mathbb{C}$, $a \in W^{1,\infty}(\mathbb{S}^{N-1}, \mathbb{R})$, $\mathbb{S}^{N-1}$ denotes the unit circle, and $\mathbf{A} \in W^{1,\infty}(\mathbb{S}^{N-1}, \mathbb{R}^N)$ is a transversal vector field, namely

$$\mathbf{A}(\theta) \cdot \theta = 0 \quad \text{for all } \theta \in \mathbb{S}^{N-1}. \tag{1.2}$$

We always denote by $r := |x|$, $\theta = x/|x|$, so that $x = r\theta$. Notice that the potentials $\mathbf{A}/|x|$ and $a/|x|^2$ preserve the natural scaling $u_\lambda(x,t) := u(x/\lambda, t/\lambda^2)$ of the free Schrödinger equation, and consequently they show a critical behavior with respect to several phenomena.

In [16], we started a program based on the connection between the Schrödinger flow $e^{it\mathcal{L}_{\mathbf{A},a}}$, generated by the hamiltonian

$$\mathcal{L}_{\mathbf{A},a} := \left(-i\nabla + \frac{\mathbf{A}\left(\frac{x}{|x|}\right)}{|x|}\right)^2 + \frac{a\left(\frac{x}{|x|}\right)}{|x|^2}, \tag{1.3}$$

and the spectral properties of the spherical operator $L_{\mathbf{A},a}$, defined by

$$L_{\mathbf{A},a} = \big(-i\,\nabla_{\mathbb{S}^{N-1}} + \mathbf{A}\big)^2 + a(\theta), \tag{1.4}$$

where $\nabla_{\mathbb{S}^{N-1}}$ is the spherical gradient on the unit sphere $\mathbb{S}^{N-1}$. In order to describe the project, let us start by reviewing some well known facts in classical spectral theory.

Date: May 7, 2014.

2000 *Mathematics Subject Classification.* 35J10, 35L05.

Key words and phrases. Schrödinger equation, electromagnetic potentials, representation formulas, decay estimates.

The first author was supported by the Italian project FIRB 2012: "Dispersive dynamics: Fourier Analysis and Variational Methods". The second author was partially supported by the P.R.I.N. 2012 grant "Variational and perturbative aspects of nonlinear differential problems". The third author was supported by the Spanish project "MTM2011-26016". The fourth author was supported by the Spanish project "MTM2010-18128".

The spectrum of the operator $L_{\mathbf{A},a}$ is formed by a diverging sequence of real eigenvalues with finite multiplicity $\mu_1(\mathbf{A},a) \leqslant \mu_2(\mathbf{A},a) \leqslant \cdots \leqslant \mu_k(\mathbf{A},a) \leqslant \cdots$ (see e.g. [18, Lemma A.5]), where each eigenvalue is repeated according to its multiplicity. Moreover we have that $\lim_{k\to\infty} \mu_k(\mathbf{A},a) = +\infty$. To each $k \geqslant 1$, we can associate a $L^2\big(\mathbb{S}^{N-1},\mathbb{C}\big)$-normalized eigenfunction ψ_k of the operator $L_{\mathbf{A},a}$ on $\mathbb{S}^{N-1}$ corresponding to the k-th eigenvalue $\mu_k(\mathbf{A},a)$, i.e. satisfying

$$\begin{cases} L_{\mathbf{A},a}\psi_k = \mu_k(\mathbf{A},a)\,\psi_k(\theta), & \text{in } \mathbb{S}^{N-1}, \\ \int_{\mathbb{S}^{N-1}} |\psi_k(\theta)|^2\, dS(\theta) = 1. \end{cases} \tag{1.5}$$

In particular, if $N = 2$, the functions ψ_k are one-variable 2π periodic functions, i.e. $\psi_k(0) = \psi_k(2\pi)$. Since the eigenvalues $\mu_k(\mathbf{A},a)$ are repeated according to their multiplicity, exactly one eigenfunction ψ_k corresponds to each index $k \geqslant 1$. We can choose the functions ψ_k in such a way that they form an orthonormal basis of $L^2(\mathbb{S}^{N-1},\mathbb{C})$. We also introduce the numbers

$$\alpha_k := \frac{N-2}{2} - \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k(\ \mathbf{A},a)}, \quad \beta_k := \sqrt{\left(\frac{N-2}{2}\right)^2 + \mu_k(\mathbf{A},a)}, \tag{1.6}$$

so that $\beta_k = \frac{N-2}{2} - \alpha_k$, for $k = 1, 2, \ldots$, which will come into play in the sequel.

Under the condition

$$\mu_1(\mathbf{A},a) > -\left(\frac{N-2}{2}\right)^2 \tag{1.7}$$

the quadratic form associated to $\mathcal{L}_{\mathbf{A},a}$ is positive definite (see [16, Section2] and [18]); this implies that the hamiltonian $\mathcal{L}_{\mathbf{A},a}$ is a symmetric semi-bounded operator on $L^2(\mathbb{R}^N;\mathbb{C})$ which then admits a self-adjoint extension (the *Friedrichs extension* which will be still denoted as $\mathcal{L}_{\mathbf{A},a}$) with domain

$$\mathcal{D}(\mathcal{L}_{\mathbf{A},a}) := \left\{ f \in H^1_*(\mathbb{R}^N) :\ \mathcal{L}_{\mathbf{A},a} u \in L^2(\mathbb{R}^N \right\}, \tag{1.8}$$

where $H^1_*(\mathbb{R}^N)$ is the completion of $C^\infty_{\rm c}(\mathbb{R}^N \setminus \{0\},\mathbb{C})$ with respect to the norm

$$\|\phi\|_{H^1_*(\mathbb{R}^N)} = \left(\int_{\mathbb{R}^N} \left(|\nabla\phi(x)|^2 + \frac{|\phi(x)|^2}{|x|^2} + |\phi(x)|^2 \right) \right) dx \Bigg)^{1/2}.$$

From the classical Hardy inequality (see e.g. [25, 28]), $H^1_*(\mathbb{R}^N) = H^1(\mathbb{R}^N)$ with equivalent norms if $N \geqslant 3$, while $H^1_*(\mathbb{R}^N)$ is strictly smaller than $H^1(\mathbb{R}^N)$ if $N = 2$. Furthermore, from condition (1.7) and [18, Lemma 2.2], it follows that $H^1_*(\mathbb{R}^N)$ coincides with the space obtained by completion of $C^\infty_{\rm c}(\mathbb{R}^N \setminus \{0\},\mathbb{C})$ with respect to the norm naturally associated to the operator $\mathcal{L}_{\mathbf{A},a}$, i.e.

$$\left(\int_{\mathbb{R}^N} \left[\left| \left(\nabla + i\, \frac{\mathbf{A}\big(x/|x|\big)}{|x|} \right) u(x) \right|^2 + \frac{a\big(x/|x|\big)}{|x|^2} |u(x)|^2 + |u(x)|^2 \right] dx \right)^{1/2}.$$

We notice that $\mathcal{L}_{\mathbf{A},a}$ could be not essentially self-adjoint. For example, in the case $\mathbf{A} \equiv 0$, from a theorem due to Kalf, Schmincke, Walter, and Wüst [30] and Simon [42] (see also [38, Theorems X.11 and X.30], [19], and [20] for non constant a), it is known that $\mathcal{L}_{\mathbf{0},a}$ is essentially self-adjoint if and only if $\mu_1(\mathbf{0},a) \geqslant -\big(\frac{N-2}{2}\big)^2 + 1$ and, consequently, admits a unique self-adjoint extension, which is given by the Friedrichs extension; otherwise, i.e. if $\mu_1(\mathbf{0},a) < -\big(\frac{N-2}{2}\big)^2 + 1$, $\mathcal{L}_{\mathbf{0},a}$ is not essentially self-adjoint and admits many self-adjoint extensions, among which the Friedrichs

extension is the only one whose domain is included in the domain of the associated quadratic form (see also [15, Remark 2.5]).

The Friedrichs extension $\mathcal{L}_{\mathbf{A},a}$ naturally extends to a self adjoint operator on the dual of $\mathcal{D}(\mathcal{L}_{\mathbf{A},a})$ and the unitary group of isometries $e^{-it\mathcal{L}_{\mathbf{A},a}}$ generated by $-i\mathcal{L}_{\mathbf{A},a}$ extends to a group of isometries on the dual of $\mathcal{D}(\mathcal{L}_{\mathbf{A},a})$ which will be still denoted as $e^{-it\mathcal{L}_{\mathbf{A},a}}$ (see [8], Section 1.6 for further details). Then for every $u_0 \in L^2(\mathbb{R}^N)$, $u(\cdot,t) = e^{-it\mathcal{L}_{\mathbf{A},a}}u_0(\cdot)$ is the unique solution to the problem

$$\begin{cases} u \in \mathcal{C}(\mathbb{R}, L^2(\mathbb{R}^N)) \cap C^1(\mathbb{R}, (\mathcal{D}(\mathcal{L}_{\mathbf{A},a}))^\star), \\ iu_t = \mathcal{L}_{\mathbf{A},a}u, \\ u(0) = u_0. \end{cases}$$

Now, by means of (1.5) and (1.6) define the following kernel:

$$K(x,y) = \sum_{k=-\infty}^{\infty} i^{-\beta_k} j_{-\alpha_k}(|x||y|)\psi_k\big(\tfrac{x}{|x|}\big)\overline{\psi_k\big(\tfrac{y}{|y|}\big)}, \tag{1.9}$$

where

$$j_\nu(r) := r^{-\frac{N-2}{2}} J_{\nu+\frac{N-2}{2}}(r)$$

and J_ν denotes the usual Bessel function of the first kind

$$J_\nu(t) = \left(\frac{t}{2}\right)^\nu \sum_{k=0}^{\infty} \frac{(-1)^k}{\Gamma(k+1)\Gamma(k+\nu+1)}\left(\frac{t}{2}\right)^{2k}.$$

In the main result of [16] we prove that, if $a \in L^\infty(\mathbb{S}^{N-1},\mathbb{R})$ and $\mathbf{A} \in C^1(\mathbb{S}^{N-1},\mathbb{R}^N)$ are such that (1.2) and (1.7) hold, then

$$e^{-it\mathcal{L}_{\mathbf{A},a}}u_0(x) = \frac{e^{\frac{i|x|^2}{4t}}}{i(2t)^{N/2}} \int_{\mathbb{R}^N} K\left(\frac{x}{\sqrt{2t}}, \frac{y}{\sqrt{2t}}\right) e^{i\frac{|y|^2}{4t}} u_0(y)\, dy, \tag{1.10}$$

for any $u_0 \in L^2(\mathbb{R}^N)$.

Apart from the interest in itself, formula (1.10) provides a quite solid tool to obtain quantitative informations for the flow $e^{-it\mathcal{L}_{\mathbf{A},a}}u_0(x)$ by the analytical study of the kernel $K(x,y)$. In particular, if

$$\sup_{x,y\in\mathbb{R}^N} |K(x,y)| < \infty \tag{1.11}$$

holds, one automatically obtains by (1.10) the time-decay estimate

$$\left\|e^{-it\mathcal{L}_{\mathbf{A},a}}u_0(\cdot)\right\|_{L^\infty} \lesssim |t|^{-\frac{N}{2}} \|u_0(\cdot)\|_{L^1}. \tag{1.12}$$

In [16], we are able to prove (1.11) (and consequently (1.12)) in two concrete situations:

- the *Aharonov-Bohm* potential: $a \equiv 0$, $\mathbf{A}(x) = \alpha\left(-\frac{x_2}{|x|}, \frac{x_1}{|x|}\right)$, for $\alpha \in \mathbb{R}$, in dimension $N = 2$;
- the positive *inverse square* potential: $\mathbf{A} \equiv 0$, $a \in \mathbb{R}$, $a > 0$.

In both cases, the spectrum of $L_{\mathbf{A},a}$ is explicit, together with a complete set of orthonormal eigenfunctions (spherical harmonics or phase transformations of themselves). These examples give a positive contribution to the recent literature about the topic, which never included before potentials with the critical homogeneity as the ones in (1.1) (see e.g. [3, 6, 7, 9, 10, 11, 12, 13, 14, 21, 22, 23, 36, 37, 39, 40, 41,

43, 44, 47, 48, 50, 51, 52, 53]). Moreover, it is well known that these potentials represent a threshold between the validity and the failure of global (in time) dispersive estimates, as proved in [17, 24]. Recently, Grillo and Kovarik [26] gave a proof of sharp time-decay estimates in the case of the Aharonov-Bohm potential, combined with a compactly supported electric potential, in dimension 2, proving also an interesting remark regarding the connection of diamagnetism with improvement of decay, in suitable weighted spaces.

The aim of this paper is to prove that estimate (1.12) holds, in space dimension $N = 2$, for a general family of potentials of the same kind as in (1.1). Our main result is the following.

Theorem 1.1. *Let $N = 2$, $a \in W^{1,\infty}(\mathbb{S}^1, \mathbb{R})$, $\mathbf{A} \in W^{1,\infty}(\mathbb{S}^1, \mathbb{R}^2)$ satisfying (1.2) and $\mu_1(\mathbf{A}, a) > 0$, and $\mathcal{L}_{\mathbf{A},a}$ be given by (1.3). Then, for any $u_0 \in L^2(\mathbb{R}^N) \cap L^1(\mathbb{R}^N)$, the following estimate holds:*

$$\left\| e^{-it\mathcal{L}_{\mathbf{A},a}} u_0(\cdot) \right\|_{L^\infty} \leqslant \frac{C}{|t|} \|u_0(\cdot)\|_{L^1}, \tag{1.13}$$

for some $C = C(\mathbf{A}, a) > 0$ which does not depend on t and u_0.

As remarked above, the proof of Theorem 1.1 consists in showing that the kernel $K(x, y)$ in (1.9) is uniformly bounded. The main difficulty is to obtain this information for the queues of the series in (1.9). In order to do this, we need to obtain the precise asymptotic behavior in k of the set of eigenvalues and eigenfunctions of the problem (1.5): this is the topic of Section 2 below. Once this is done, the proof of Theorem 1.1 will be obtained, in Section 3, by suitably comparing the kernel K with the analogous in the case of an Aharonov-Bohm potential with the same average as the potential $\mathbf{A}$ on the sphere $\mathbb{S}^1$.

Acknowledgements. The authors wish to thank Carlos Villegas-Blas for addressing them to several useful references about the topic of Section 2.

2. Spectral properties of spherical laplacians

The fundamental tool which we need in order to prove Theorem 1.1 is the knowledge of the spectral properties of the operator $L_{\mathbf{A},a}$ defined by (1.4). Roughly speaking, we need to obtain informations concerning the asymptotic behavior of the eigenvalues μ_k and the eigenfunctions ψ_k in the eigenvalue problem (1.5), as $k \to \infty$.

An extensive literature has been devoted, in the recent years, to this kind of problems (see e.g. [27, 45, 46, 49] and the references therein). Since we did not find sufficiently explicit results regarding general electromagnetic Laplace operators on the 1D-sphere $\mathbb{S}^1$, we need to show here Lemma 2.1 below, which is possibly of independent interest.

Before starting to settle the eigenvalue problem, we find convenient to briefly sketch the well known consequences which the introduction of lower order terms produces on the spectrum of the spherical Laplacian.

Let us denote by $L_0 := -\Delta_{\mathbb{S}^1}$. Being the inverse of a compact operator on $L^2(\mathbb{S}^1)$, with form domain $H^1(\mathbb{S}^1)$, L_0 has purely discrete spectrum which accumulates at positive infinity. The explicit form is

$$\sigma(L_0) = \{k^2\}_{k=0,1,\ldots}.$$

The k^{th}-eigenvalue has multiplicity 2, and the eigenfunctions are combinations of sines and cosines.

The introduction of a 0-order term produces a spectral shift, depending on the average of the potentials, and the formation of clusters of eigenvalues around the free ones (Stark's effect), if the potential is not constant. More precisely, the eigenvalues of the operator $L_a := -\Delta_{\mathbb{S}^1} + a(\theta)$ are of the form

$$\lambda_k = k^2 + \frac{1}{2\pi}\int_0^{2\pi} a(s)\,ds + (\text{rest}),$$

where the rest, depending on k and on the potential a, decays with order $1/k$ as k tends to infinity. For the eigenfunctions ψ_k a similar behavior occurs; for large k, ψ_k looks more and more like a spherical harmonic plus a rest which decays as k tends to $+\infty$ (see e.g. [5, 27, 30, 45, 46] and appendix B).

On the other hand, for a purely magnetic potential, a splitting occurs on each eigenvalue. The most famous (and descriptive) example is given by the Aharonov-Bohm potential, namely $a \equiv 0$, $\mathbf{A}(x,y) = \mathbf{A}_{ab}(x,y) = \alpha\left(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2}\right)$, with $\alpha \in \mathbb{R}$: in this case, the complete set of eigenvalues and eigenfunctions of problem (1.5) can be computed explicitly, and reads as

$$\lambda_k^{ab} = (k+\alpha)^2, \quad k \in \mathbb{Z} \tag{2.1}$$

$$\phi_k^{ab}(\theta) = \frac{1}{\sqrt{2\pi}} e^{ik\theta}, \quad k \in \mathbb{Z}. \tag{2.2}$$

It is hence quite natural to expect that, in the general case of the operator $L_{\mathbf{A},a}$, the picture is a superposition of the two previously mentioned ones. We did not find in the literature a result written in the generality of Lemma 2.1 below, so that we found convenient to state and prove it in this manuscript.

We recall that, by classical spectral theory, the spectrum of $L_{\mathbf{A},a}$ is formed by a countable family of real eigenvalues with finite multiplicity $\{\mu_k : k \geqslant 1\}$ enumerated in such a way that

$$\mu_1 \leqslant \mu_2 \leqslant \dots,$$

where each eigenvalue is repeated according to its multiplicity. Moreover we have that $\lim_{k\to\infty} \mu_k = +\infty$.

Let $A : [0, 2\pi] \to \mathbb{R}$ be defined as $A(\theta) = \mathbf{A}(\cos\theta, \sin\theta) \cdot (-\sin\theta, \cos\theta)$, so that, by assumption (1.2)

$$\mathbf{A}(\cos\theta, \sin\theta) = A(\theta)(-\sin\theta, \cos\theta), \quad \theta \in [0, 2\pi]. \tag{2.3}$$

Furthermore, identifying functions defined on $\mathbb{S}^1$ with 2π-periodic functions, the operator $L_{\mathbf{A},a}$ can be identified with the following operator $\mathfrak{L}_{A,a}$ acting on 2π-periodic functions

$$\mathfrak{L}_{A,a}\varphi(\theta) = -\varphi''(\theta) + [a(\theta) + A^2(\theta) - iA'(\theta)]\varphi(\theta) - 2iA(\theta)\varphi'(\theta).$$

The main result of this section is the following asymptotic expansion of eigenvalues and eigenfunctions of the operator $L_{\mathbf{A},a}$ under the non-resonant assumption that the magnetic potential does not have half-integer or integer circulation. The case of half-integer or integer circulation can be reduced through suitable transformations to the magnetic-free problem, for which analogous expansions hold, see Remark 2.2 and appendix B.

Lemma 2.1. *Let $a \in W^{1,\infty}(\mathbb{S}^1)$, $\widetilde{a} = \frac{1}{2\pi}\int_0^{2\pi} a(s)\,ds$, $A \in W^{1,\infty}(\mathbb{S}^1)$ such that*

$$\widetilde{A} = \frac{1}{2\pi}\int_0^{2\pi} A(s)\,ds \notin \frac{1}{2}\mathbb{Z}. \tag{2.4}$$

Then there exist $k^*, \ell \in \mathbb{N}$ such that $\{\mu_k :\ k > k^*\} = \{\lambda_j :\ j \in \mathbb{Z},\ |j| \geqslant \ell\}$,

$$\sqrt{\lambda_j - \widetilde{a}} = (\operatorname{sgn} j)\Big(\widetilde{A} - \Big\lfloor \widetilde{A} + \tfrac{1}{2}\Big\rfloor\Big) + |j| + O\big(\tfrac{1}{|j|^3}\big), \quad \text{as } |j| \to +\infty$$

and

$$\lambda_j = \widetilde{a} + \Big(j + \widetilde{A} - \big\lfloor \widetilde{A} + \tfrac{1}{2}\big\rfloor\Big)^2 + O\big(\tfrac{1}{j^2}\big), \quad \text{as } |j| \to +\infty. \tag{2.5}$$

Furthermore, for all $j \in \mathbb{Z}$, $|j| \geqslant \ell$, there exists a $L^2\big(\mathbb{S}^1, \mathbb{C}\big)$-normalized eigenfunction ϕ_j of the operator $L_{\mathbf{A},a}$ on $\mathbb{S}^1$ corresponding to the eigenvalue λ_j such that

$$\phi_j(\theta) = \frac{1}{\sqrt{2\pi}} e^{-i\left([\widetilde{A}+1/2]\theta + \int_0^\theta A(t)\,dt\right)}\Big(e^{i(\widetilde{A}+j)\theta} + R_j(\theta)\Big), \tag{2.6}$$

where $\|R_j\|_{L^\infty(\mathbb{S}^1)} = O\big(\frac{1}{|j|^3}\big)$ as $|j| \to \infty$. In the above formula $\lfloor\cdot\rfloor$ denotes the floor function $\lfloor x \rfloor = \max\{k \in \mathbb{Z} : k \leqslant x\}$.

Lemma 2.1 can be interpreted as follows: asymptotically in k, eigenvalues and eigenfunctions of (1.5) for $L_{\mathbf{A},a}$ are comparable with the ones in the Aharonov-Bohm case (see (2.1), (2.2) above), by means of (2.5), (2.6).

The proof of Lemma 2.1 is based on the idea of reducing the eigenvalue problem (1.5) to another magnetic-free problem, with different boundary conditions, by gauge transformation; this is in fact possible, since $\mathbf{A}(\cos\theta, \sin\theta)$ just depends on the 1D-variable θ. More precisely, we observe that the gauge transformation

$$\psi(\theta) \to e^{-i\int_0^\theta A(s)\,ds}\psi(\theta)$$

transforms the eigenvalue problem (1.5) into the new problem

$$\begin{cases} -\frac{d^2\psi}{d\theta^2} + a(\theta)\psi = \mu_k\psi \\ \psi(0) = e^{-2i\pi\widetilde{A}}\psi(2\pi) \\ \psi'(0) = e^{-2i\pi\widetilde{A}}\psi'(2\pi), \end{cases} \tag{2.7}$$

with non-periodic boundary conditions, where $\widetilde{A}$ is defined in (2.4), which will be analyzed by a usual WKB-strategy.

Remark 2.2. As mentioned above, in the purely electric case $\mathbf{A} \equiv 0$, (2.5) is a well known information about the cluster distribution of the eigenvalues (see e.g. [27] and the references therein). More in general, if $\operatorname{dist}(\widetilde{A}, \mathbb{Z}) = 0$, then the eigenvalue problem (2.7) reduces to

$$\begin{cases} -\frac{d^2\psi_k}{d\theta^2} + a(\theta)\psi_k = \mu_k\psi_k \\ \psi_k(0) = \psi_k(2\pi) \\ \psi_k'(0) = \psi_k'(2\pi), \end{cases} \tag{2.8}$$

i.e. the magnetic-free case. For the proof of Lemma 2.1 in the case $\operatorname{dist}(\widetilde{A}, \mathbb{Z}) = 0$ we mention a vintage result by Borg [5] (see also [27]) as a standard reference; for the sake of completeness we sketch a proof of asymptotics of eigenvalues and eigenfunctions in the purely electric case in the appendix. Nevertheless, we propose here a proof in the case $\operatorname{dist}(\widetilde{A}, \mathbb{Z}) \neq 0, \frac{1}{2}$, since we did not find in the literature

neither the analogous to [5] for $\mathbf{A} \neq 0$ nor the asymptotic formula for eigenfunctions (2.6), which plays a fundamental role in the proof of our main theorem (see section 3 below). We propose a proof which is based on a usual WKB-strategy.

2.1. **Proof of Lemma 2.1.** Let us denote

$$\bar{A} = \widetilde{A} - \left\lfloor \widetilde{A} + \frac{1}{2} \right\rfloor, \tag{2.9}$$

so that $\bar{A} \in [-1/2, 1/2)$; we notice that $\widetilde{A} \in \frac{1}{2}\mathbb{Z}$ if and only if $\bar{A} \in \{-1/2, 0\}$. Hence, under assumption (2.4), we have that

$$\bar{A} \in \Big(-\frac{1}{2}, \frac{1}{2} \Big) \setminus \{0\}. \tag{2.10}$$

Lemma 2.3. *Let $a, A \in W^{1,\infty}(0, 2\pi)$, $\widetilde{A} = \frac{1}{2\pi}\int_0^{2\pi} A(s)\,ds$, and $\bar{A}$ as in (2.9), i.e. $\bar{A} = \widetilde{A} - \lfloor \widetilde{A} + \frac{1}{2} \rfloor$. Then, letting $\mathbf{A}$ as in (2.3), we have that*

$$\sigma(L_{\mathbf{A},a}) = \sigma(\mathfrak{L}_{A,a}) = \sigma(\mathfrak{L}_{\bar{A},a}).$$

Furthermore, φ is an eigenfunction of $\mathfrak{L}_{A,a}$ associated to the eigenvalue μ if and only if $\widetilde{\varphi}(t) = e^{-i\bar{A}t} e^{i\int_0^t A(s)\,ds}\varphi(t)$ is an eigenfunction of $\mathfrak{L}_{\bar{A},a}$ associated to μ.

Proof. The proof follows by direct calculations. We notice that, since $\widetilde{A} - \bar{A} \in \mathbb{Z}$, function $\widetilde{\varphi}(t) = e^{-i\bar{A}t} e^{i\int_0^t A(s)\,ds}\varphi(t)$ is 2π-periodic if and only if $\varphi(t)$ is 2π-periodic. □

Lemma 2.4. *Let $a \in W^{1,\infty}(\mathbb{S}^1)$, $\widetilde{a} = \frac{1}{2\pi}\int_0^{2\pi} a(s)\,ds$, $\delta > 0$, and*

$$I_\delta = \{\lambda \in \mathbb{R} : \ \operatorname{dist}(\sqrt{\lambda - \widetilde{a}}, \tfrac{1}{2}\mathbb{Z}) \geqslant \delta\}.$$

There exist $\bar{\lambda}_\delta > 0$ and $C_\delta > 0$ such that for every $\lambda \in I_\delta$, $\lambda \geqslant \bar{\lambda}_\delta$, there exists $W_\lambda \in C^0(\mathbb{S}^1)$ such that

$$\|W_\lambda\|_{C^0(\mathbb{S}^1)} \leqslant \frac{C_\delta}{\sqrt{\lambda - \widetilde{a}}} \tag{2.11}$$

and

$$T_\lambda(W_\lambda) = W_\lambda,$$

where $T_\lambda : C^0(\mathbb{S}^1) \to C^0(\mathbb{S}^1)$ is defined as

$$\begin{aligned}
T_\lambda(W)(\theta) &= e^{-2\sqrt{\lambda-\widetilde{a}}\theta i}\frac{\widetilde{a}-a(0)}{2\sqrt{\lambda-\widetilde{a}}} \\
&\quad + \frac{ie^{-2\sqrt{\lambda-\widetilde{a}}(\theta+2\pi)i}}{1-e^{-4\sqrt{\lambda-\widetilde{a}}\pi i}}\int_0^{2\pi} e^{2\sqrt{\lambda-\widetilde{a}}\theta' i}\left(\frac{a'(\theta')}{2\sqrt{\lambda-\widetilde{a}}i} - W^2(\theta')\right)d\theta' \\
&\quad + ie^{-2\sqrt{\lambda-\widetilde{a}}\theta i}\int_0^{\theta} e^{2\sqrt{\lambda-\widetilde{a}}\theta' i}\left[\widetilde{a}-a(\theta')-W^2(\theta')\right]d\theta' \\
&= e^{-2\sqrt{\lambda-\widetilde{a}}\theta i}\frac{\widetilde{a}-a(0)}{2\sqrt{\lambda-\widetilde{a}}} \\
&\quad + \frac{ie^{-2\sqrt{\lambda-\widetilde{a}}(\theta+2\pi)i}}{1-e^{-4\sqrt{\lambda-\widetilde{a}}\pi i}}\int_0^{2\pi} e^{2\sqrt{\lambda-\widetilde{a}}\theta' i}\left(\frac{a'(\theta')}{2\sqrt{\lambda-\widetilde{a}}i} - W^2(\theta')\right)d\theta' \\
&\quad - \frac{a(\theta)-e^{-2\sqrt{\lambda-\widetilde{a}}\theta i}a(0)}{2\sqrt{\lambda-\widetilde{a}}} + \frac{\widetilde{a}-e^{-2\sqrt{\lambda-\widetilde{a}}\theta i}\widetilde{a}}{2\sqrt{\lambda-\widetilde{a}}} \\
&\quad + ie^{-2\sqrt{\lambda-\widetilde{a}}\theta i}\int_0^{\theta} e^{2\sqrt{\lambda-\widetilde{a}}\theta' i}\left[\frac{a'(\theta')}{2\sqrt{\lambda-\widetilde{a}}i} - W^2(\theta')\right]d\theta'.
\end{aligned}$$

Moreover the map $\lambda \mapsto W_\lambda$ is continuous as a map from I_δ to $C^0(\mathbb{S}^1)$.

Proof. It is easy to verify that there exist $\bar\lambda_\delta$ and $C_\delta > 0$ such that for every $\lambda \in I_\delta$, $\lambda \geqslant \bar\lambda_\delta$, T_λ maps $\overline{B}_{C_\delta/\sqrt{\lambda-\widetilde{a}}} = \{u \in C^0(\mathbb{S}^1) : \sup_{\mathbb{S}^1}|u| \leqslant C_\delta/\sqrt{\lambda-\widetilde{a}}\}$ into itself and is a contraction there. The conclusion then follows from the Banach contraction mapping theorem. □

For $\lambda \in I_\delta$, $\lambda \geqslant \bar\lambda_\delta$, let W_λ be as in Lemma 2.4. Then it is easy to verify that W_λ satisfies

$$\begin{cases} -iW_\lambda'(\theta) + 2\sqrt{\lambda-\widetilde{a}}\,W_\lambda(\theta) + W_\lambda^2(\theta) = \widetilde{a}-a(\theta), \quad \text{in } [0,2\pi], \\ W_\lambda(0) = W_\lambda(2\pi). \end{cases} \tag{2.12}$$

Letting

$$S_\lambda(\theta) := \sqrt{\lambda-\widetilde{a}}\,\theta + \int_0^\theta W_\lambda(\theta')\,d\theta', \tag{2.13}$$

we have that S_λ satisfies

$$\begin{cases} -iS_\lambda''(\theta) + (S_\lambda'(\theta))^2 = \lambda - a(\theta), \quad \text{in } [0,2\pi], \\ S_\lambda'(0) = S_\lambda'(2\pi), \\ S_\lambda(0) = 0. \end{cases} \tag{2.14}$$

Lemma 2.5. *If λ is sufficiently large, then $\int_0^{2\pi} W_\lambda(\theta)\,d\theta \in \mathbb{R}$.*

Proof. Let us define $\eta_\lambda(\theta) = \Re S_\lambda(\theta)$ and $\xi_\lambda(\theta) = \Im S_\lambda(\theta)$. Then (2.14) implies that

$$-\eta_\lambda'' + 2\eta_\lambda'\xi_\lambda' = 0, \quad \text{in } [0,2\pi],$$

so that

$$\eta_\lambda'(\theta) = C_\lambda e^{2\xi_\lambda(\theta)}, \quad \text{in } [0,2\pi], \tag{2.15}$$

where $C_\lambda = \sqrt{\lambda - \widetilde{a}} + \Re W_\lambda(0)$. We notice that (2.11) implies that, if λ is sufficiently large, then $C_\lambda \neq 0$. The condition $S'_\lambda(0) = S'_\lambda(2\pi)$ implies that $\eta'_\lambda(0) = \eta'_\lambda(2\pi)$ and hence from (2.15) it follows that

$$\xi_\lambda(0) = \xi_\lambda(2\pi).$$

Since $S_\lambda(0) = 0$, we have that $\xi_\lambda(0) = 0$ and then

$$\begin{aligned}\frac{1}{2\pi}\int_0^{2\pi} W_\lambda(\theta)\,d\theta &= -\sqrt{\lambda - \widetilde{a}} + \frac{1}{2\pi}(\eta_\lambda(2\pi) + i\xi_\lambda(2\pi)) \\ &= -\sqrt{\lambda - \widetilde{a}} + \frac{\eta_\lambda(2\pi)}{2\pi} \in \mathbb{R}.\end{aligned}$$

□

Lemma 2.6. *Let $\bar{A} \in \mathbb{R}$ such that $\bar{A} \notin \frac{1}{2}\mathbb{Z}$ and let $0 < \delta < \operatorname{dist}(\bar{A}, \frac{1}{2}\mathbb{Z})$. Then there exists $\bar{k} \in \mathbb{N}$ such that for all $k \in \mathbb{N}$, $k \geqslant \bar{k}$, there exist $\lambda_k^+, \lambda_k^- \in I_\delta$ such that $\lambda_k^+ \geqslant \bar{\lambda}_\delta$, $\lambda_k^- \geqslant \bar{\lambda}_\delta$ and*

$$\sqrt{\lambda_k^+ - \widetilde{a}} = \bar{A} - \frac{1}{2\pi}\int_0^{2\pi} W_{\lambda_k^+}(\theta)\,d\theta + k, \tag{2.16}$$

$$\sqrt{\lambda_k^- - \widetilde{a}} = -\bar{A} - \frac{1}{2\pi}\int_0^{2\pi} W_{\lambda_k^-}(\theta)\,d\theta + k. \tag{2.17}$$

Proof. Let $g : [\bar{\lambda}_\delta, +\infty) \to \mathbb{R}$ be a continuous function such that

$$g(\lambda) = \frac{1}{2\pi}\int_0^{2\pi} W_\lambda(\theta)\,d\theta \quad \text{for all } \lambda \in I_\delta$$

and $|g(\lambda)| \leqslant C_\delta/\sqrt{\lambda - \tilde{a}}$ for all $\lambda \geqslant \bar{\lambda}_\delta$. Then the function

$$f : [\bar{\lambda}_\delta, +\infty) \to \mathbb{R}, \quad f(\lambda) = \sqrt{\lambda - \tilde{a}} - \bar{A} + g(\lambda),$$

is continuous and $\lim_{\lambda\to+\infty} f(\lambda) = +\infty$. Therefore there exists $\bar{k}$ sufficiently large such that, for all $k \geqslant \bar{k}$, there exists $\lambda_k^+ \geqslant \bar{\lambda}_\delta$ such that $f(\lambda_k^+) = k$, i.e.

$$\sqrt{\lambda_k^+ - \tilde{a}} = k + \bar{A} - g(\lambda_k^+). \tag{2.18}$$

If $\bar{k}$ is sufficiently large, (2.18) implies that

$$\operatorname{dist}\left(\sqrt{\lambda_k^+ - \tilde{a}}, \tfrac{1}{2}\mathbb{Z}\right) = \operatorname{dist}(\bar{A}, \tfrac{1}{2}\mathbb{Z}) - g(\lambda_k^+) > \delta,$$

so that $\lambda_k^+ \in I_\delta$ and (2.16) is proved. The proof of (2.17) is analogous. □

Lemma 2.7. *Under the same assumptions as in Lemma 2.6, let, for all $k \geqslant \bar{k}$, $\lambda_k^+, \lambda_k^- \in I_\delta$ as in Lemma 2.6. Then*

$$\int_0^{2\pi} W_{\lambda_k^\pm}(\theta)\,d\theta = O\big(\tfrac{1}{k^3}\big), \quad \text{as } k \to +\infty \tag{2.19}$$

and

$$\sqrt{\lambda_k^+ - \widetilde{a}} = \bar{A} + k + O(\tfrac{1}{k^3}), \quad \sqrt{\lambda_k^- - \widetilde{a}} = -\bar{A} + k + O(\tfrac{1}{k^3}), \tag{2.20}$$

$$\lambda_k^+ = \widetilde{a} + (\bar{A} + k)^2 + O(\tfrac{1}{k^2}), \quad \lambda_k^- = \widetilde{a} + (-\bar{A} + k)^2 + O(\tfrac{1}{k^2}), \tag{2.21}$$

as $k \to +\infty$.

PROOF. By integrating (2.12) between 0 and 2π and using estimate (2.11) we have that

$$\left| \int_0^{2\pi} W_\lambda(\theta)\, d\theta \right| = \left| -\frac{1}{2\sqrt{\lambda - \widetilde{a}}} \int_0^{2\pi} W_\lambda^2(\theta)\, d\theta \right| \leqslant \frac{\pi C_\delta}{(\lambda - \widetilde{a})^{3/2}}.$$

Since from (2.16) and (2.17) it follows that $\lambda_k^\pm \sim k^2$ as $k \to +\infty$, we derive (2.19), which yields (2.20) (and the (2.21) by squaring) in view of (2.16) and (2.17). □

Lemma 2.8. *Let $a \in W^{1,\infty}(\mathbb{S}^1)$ and $\bar{A} \in \mathbb{R} \setminus \frac{1}{2}\mathbb{Z}$. If $k \geqslant \bar{k}$, then*

$$\lambda_k^+, \lambda_k^- \in \sigma(\mathfrak{L}_{\bar{A},a}),$$

where $\mathfrak{L}_{\bar{A},a}\varphi = -(\varphi)'' + [a(\theta) + \bar{A}^2]\varphi - 2i\bar{A}\varphi$. Moreover

$$\varphi_k^+(\theta) = e^{-i\bar{A}\theta} e^{iS_{\lambda_k^+}(\theta)}, \quad \varphi_k^-(\theta) = e^{-i\bar{A}\theta} e^{-i\overline{S_{\lambda_k^-}(\theta)}}, \tag{2.22}$$

are eigenfunctions of $\mathfrak{L}_{\bar{A},a}$ associated to λ_k^+, λ_k^- respectively.

PROOF. By direct calculations, we have that $\varphi_k^\pm$ satisfy

$$-(\varphi_k^\pm)''(\theta) + [a(\theta) + \bar{A}^2]\varphi_k^\pm(\theta) - 2i\bar{A}(\varphi_k^\pm)'(\theta) = \lambda_k^\pm \varphi_k^\pm(\theta)$$

in $[0, 2\pi]$, i.e. $\varphi_k^\pm$ are non-trivial solutions to

$$\mathfrak{L}_{\bar{A},a}\varphi_k^\pm = \lambda_k^\pm \varphi_k^\pm \quad \text{in } [0, 2\pi].$$

Furthermore (2.16) and (2.17) imply that

$$\varphi_k^\pm(0) = \varphi_k^\pm(2\pi) \quad \text{and} \quad (\varphi_k^\pm)'(0) = (\varphi_k^\pm)'(2\pi).$$

The lemma is thereby proved. □

Lemma 2.9. *Let $a \in W^{1,\infty}(\mathbb{S}^1)$ and $\bar{A} \in (-\frac{1}{2}, \frac{1}{2}) \setminus \{0\}$. For every $\bar{\alpha} \geqslant \|a + \bar{A}^2\|_{L^\infty}^2$ and $c \geqslant 0$, there exist $\bar{\lambda} > 0$ and $k_0 > \bar{k}$ such that*

$$\sigma(\mathfrak{L}_{\bar{A},a}) \cap [\bar{\lambda}, +\infty) \subset \bigcup_{k=k_0}^{\infty} B\big(k^2, c + \sqrt{\bar{\alpha} + 4k^2\bar{A}^2}\big).$$

Furthermore, if $k \geqslant k_0$, each ball $B\big(k^2, \sqrt{\bar{\alpha} + 4k^2A^2}\big)$ contains exactly two eigenvalues of $\mathfrak{L}_{\bar{A},a}$ (counted with their own multiplicity).

PROOF. We apply lemma A.1 with

$$L_0 := -\frac{d^2}{d\theta^2} \qquad L := -\frac{d^2}{d\theta^2} - 2i\bar{A}\frac{d}{d\theta} + \alpha(\theta),$$

where $\alpha(\theta) = a(\theta) + \bar{A}^2$. Let

$$R_0(\lambda) = \Big(-\frac{d^2}{d\theta^2} - \lambda I \Big)^{-1}, \quad R(\lambda) = \Big(-\frac{d^2}{d\theta} - 2i\bar{A}\frac{d}{d\theta} + \alpha(\theta) - \lambda I \Big)^{-1}.$$

Notice that, via Fourier we can write, for $f = \sum(\alpha_k \sin(k\theta) + \beta_k \cos(k\theta))$,

$$R_0(\lambda) f = \sum \frac{1}{k^2 - \lambda}(\alpha_k \sin(k\theta) + \beta_k \cos(k\theta)),$$

therefore we have the estimate

$$\|R_0\|_{\mathcal{L}(L^2)} \leqslant \frac{1}{\operatorname{dist}(\lambda, \{k^2 : k \in \mathbb{Z}\})}.$$

Now notice that, formally, we can write

$$R(\lambda) = R_0(\lambda)\left(I + WR_0(\lambda)\right)^{-1}, \tag{2.23}$$

being $W = -2i\bar{A}\frac{d}{d\theta} + \alpha(\theta)$ a first order operator. Since $\frac{d}{d\theta}$ commutes with R_0 (by spectral theorem), we can write as follows:

$$WR_0(\lambda)f = -2i\bar{A}\sum_k \frac{k(\alpha_k\cos(k\theta) - \beta_k\sin(k\theta))}{k^2-\lambda} + \alpha(\theta)\sum_k \frac{1}{k^2-\lambda}(\alpha_k\sin(k\theta) + \beta_k\cos(k\theta)).$$

We hence obtain the following: if $\Re\lambda$ is large enough, $\bar{\alpha} \geqslant \|\alpha\|^2_{L^\infty}$, $c \geqslant 0$, and

$$|\lambda - k^2| > c + \left(4k^2\bar{A}^2 + \bar{\alpha}\right)^{\frac{1}{2}},$$

then

$$\|WR_0\|_{\mathcal{L}(L^2)} < 1. \tag{2.24}$$

Then, by (2.23) and (2.24), if k is large enough, outside of any ball with center in k^2 and radius $c + \left(4k^2\bar{A}^2 + \bar{\alpha}\right)^{\frac{1}{2}}$, the operator $R(\lambda)$ is bounded for large λ, hence we do not have large eigenvalues outside that balls. We notice that, if k is large, the balls with center in k^2 and radius $c + \left(4k^2\bar{A}^2 + \bar{\alpha}\right)^{\frac{1}{2}}$ are mutually disjoint, since $|\bar{A}| < \frac{1}{2}$ implies that

$$2c + \left(4k^2\bar{A}^2 + \bar{\alpha}\right)^{\frac{1}{2}} + \left(4(k+1)^2\bar{A}^2 + \bar{\alpha}\right)^{\frac{1}{2}} < (k+1)^2 - k^2$$

provided that k is sufficiently large. On the other hand, if Γ is the circle with center in k^2 and radius $\left(4k^2\bar{A}^2 + \bar{\alpha}\right)^{\frac{1}{2}}$ with k large, we can easily estimate

$$\left\| \frac{1}{2\pi i}\int_\Gamma (R(\lambda) - R_0(\lambda))f\, d\lambda \right\|_{L^2} < \|f\|_{L^2},$$

(use the Born expansion $(I + WR_0)^{-1} = \sum (WR_0)^n$) which together with point (2) of Lemma A.1 gives the desired result.

Therefore, outside those balls there are no eigenvalues, and inside there are the same number of eigenvalues both for L_0 and L: this number is 2. □

Lemma 2.10. *Let $a \in W^{1,\infty}(\mathbb{S}^1)$ and $\bar{A} \in (-\frac{1}{2}, \frac{1}{2}) \setminus \{0\}$. Let $\lambda_k^\pm$ be as in Lemma 2.6. Then there exist $c > 0$, $\bar{\alpha} > 0$, $\bar{\lambda} > 0$ and $\tilde{k}$ such that*

(i) *for all $k \geqslant \tilde{k}$, $\lambda_k^+, \lambda_k^- \in B\left(k^2, c + \sqrt{\bar{\alpha} + 4k^2\bar{A}^2}\right)$;*

(ii) $\sigma(\mathfrak{L}_{\bar{A},a}) \cap [\tilde{\lambda}, +\infty) = \{\lambda_k^+, \lambda_k^- : k \geqslant \tilde{k}\}$.

Proof. From (2.21) we have that, if $c, \bar{\alpha} > 0$ are chosen sufficiently large,

$$|\lambda_k^+ - k^2| = \left|\widetilde{a} + \bar{A}^2 + 2k\bar{A} + O(\tfrac{1}{k^2})\right| < c + \sqrt{\bar{\alpha} + 4k^2\bar{A}^2}$$

if k is large enough, thus proving (i) for λ_k^+. The proof of (i) for λ_k^- is analogous.

The statement (ii) follows by combining (i) and Lemma 2.9. □

Proof of Lemma 2.1. From Lemmas 2.3 and 2.10 it follows that there exist $k^* \in \mathbb{N}$ and $\ell \in \mathbb{Z}$ such that $\{\mu_k : \ k > k^*\} = \{\lambda_j : \ j \in \mathbb{Z}, \ |j| \geqslant \ell\}$ where

$$\lambda_j = \begin{cases} \lambda^-_{|j|}, & \text{if } j < 0, \\ \lambda^+_{|j|}, & \text{if } j > 0. \end{cases}$$

Then, in view of Lemma 2.7

$$\sqrt{\lambda_j - \widetilde{a}} = (\operatorname{sgn} j)\bar{A} + |j| + O\big(\tfrac{1}{|j|^3}\big), \quad \text{as } |j| \to +\infty.$$

From (2.22), (2.13), (2.19), and (2.20), it follows that

$$\varphi^+_j(\theta) = e^{-i\bar{A}\theta}\Big(e^{i(\bar{A}+j)\theta} + O\big(\tfrac{1}{|j|^3}\big)\Big), \quad \text{as } j \to +\infty,$$
$$\varphi^-_j(\theta) = e^{-i\bar{A}\theta}\Big(e^{i(\bar{A}-j)\theta} + O\big(\tfrac{1}{|j|^3}\big)\Big), \quad \text{as } j \to +\infty.$$

Therefore, letting, for $j \in \mathbb{Z}$ such that $|j| \geqslant \ell$,

$$\widetilde{\phi}_j = \begin{cases} \frac{\varphi^-_{|j|}}{\|\varphi^-_{|j|}\|_{L^2(0,2\pi)}}, & \text{if } j < 0, \\ \frac{\varphi^+_{|j|}}{\|\varphi^+_{|j|}\|_{L^2(0,2\pi)}}, & \text{if } j > 0, \end{cases}$$

we have that, for $|j| \geqslant \ell$, $\widetilde{\phi}_j$ is a $L^2\big((0,2\pi),\mathbb{C}\big)$-normalized eigenfunction of the operator $\mathfrak{L}_{\bar{A},a}$ corresponding to the eigenvalue λ_j and

$$\widetilde{\phi}_j(\theta) = \frac{1}{\sqrt{2\pi}} e^{-i\bar{A}\theta}\Big(e^{i(\bar{A}+j)\theta} + R_j(\theta)\Big),$$

where $\|R_j\|_{L^\infty(0,2\pi)} = O\big(\frac{1}{|j|^3}\big)$ as $j \to \infty$. Hence, in view of Lemma 2.3 we have that $\phi_j(\cos\theta, \sin\theta) = e^{i\bar{A}\theta} e^{-i\int_0^\theta A(s)\,ds}\widetilde{\phi}_j(\theta)$ is a $L^2\big(\mathbb{S}^1,\mathbb{C}\big)$-normalized eigenfunction of the operator $L_{\mathbf{A},a}$ on $\mathbb{S}^1$ corresponding to the eigenvalue λ_j and

$$\phi_j(\cos\theta, \sin\theta) = \frac{1}{\sqrt{2\pi}} e^{-i\left(\lfloor \widetilde{A}+1/2\rfloor\theta + \int_0^\theta A(t)\,dt\right)}\Big(e^{i(\widetilde{A}+j)\theta} + R_j(\theta)\Big).$$

The proof is thereby complete. □

By means of the previous result, we immediately obtain the following Corollary.

Corollary 2.11. *Let k^*, ℓ as in Lemma 2.1 and K be given by* (1.9)*, with ψ_k being any $L^2\big(\mathbb{S}^1,\mathbb{C}\big)$-normalized eigenfunctions of $L_{\mathbf{A},a}$ on $\mathbb{S}^1$ if $k \leqslant k^*$ and $\psi_k = \phi_j$ if $k > k^*$ and $\mu_k = \lambda_j$, with λ_j, ϕ_j being as in Lemma 2.1. Then, we have that*

$$\begin{aligned} K(x,y) = &\sum_{k=1}^{k^*} i^{-\beta_k} j_{-\alpha_k}(rr')\psi_k(\theta)\overline{\psi_k(\theta')} \\ &+ \frac{1}{2\pi} e^{-i\int_{\theta'}^{\theta} A(s)\,ds} e^{-i[\widetilde{A}+\frac{1}{2}](\theta-\theta')} \\ &\quad \times \sum_{|j|\geqslant\ell} i^{-\beta(\lambda_j)} j_{-\alpha(\lambda_j)}(rr')\Big(e^{i(\widetilde{A}+j)\theta} + R_j(\theta)\Big)\Big(e^{-i(\widetilde{A}+j)\theta'} + \overline{R_j(\theta')}\Big), \end{aligned} \tag{2.25}$$

if $x = (r\cos\theta, r\sin\theta)$ and $y = (r'\cos\theta', r'\sin\theta')$, where

$$\alpha(\lambda_j) := -\sqrt{\lambda_j}, \quad \beta(\lambda_j) := \sqrt{\lambda_j}, \tag{2.26}$$

and R_j is as in Lemma 2.1.

3. Proof of the main result

We can now perform the proof of Theorem 1.1. Let us first assume that condition (2.4) holds, so that the asymptotic expansion of eigenvalues and eigenfunctions stated in Lemma 2.1 holds. Let K be defined by (1.9); by formula (1.10), it is sufficient to show that

$$\sup_{x,y\in\mathbb{R}^2} |K(x,y)| < \infty.$$

In particular, the study of the boundedness of K is reduced, thanks to Corollary 2.11, to the study of the boundedness of the two series

$$\Sigma_{k\leqslant k^*} = \sum_{k=1}^{k^*} i^{-\beta_k} j_{-\alpha_k}(rr')\psi_k(\theta)\overline{\psi_k(\theta')}, \tag{3.1}$$

and

$$\Sigma_{|j|\geqslant\ell} = \sum_{|j|\geqslant\ell} i^{-\beta(\lambda_j)} j_{-\alpha(\lambda_j)}(rr')\Big(e^{i(\widetilde{A}+j)\theta} + R_j(\theta)\Big)\Big(e^{-i(\widetilde{A}+j)\theta'} + \overline{R_j(\theta')}\Big) \tag{3.2}$$

uniformly with respect to r, r', θ, θ'. Since $\mu_1(\mathbf{A}, a) > 0$, all the indices α_k in (1.6) are negative. Therefore, the Bessel functions $j_{-\alpha_k}$ are bounded functions, for any k. In addition, the functions ψ_k are obviously bounded, for any k: as a consequence, we obtain that

$$\sup_{\substack{r,r'\geqslant 0\\ \theta,\theta'\in\mathbb{S}^1}} |\Sigma_{k\leqslant k^*}(r,r',\theta,\theta')| < \infty. \tag{3.3}$$

In order to prove that $\Sigma_{|j|\geqslant\ell}$ is uniformly bounded, we compare it with the analogous kernel K_{ab} associated to the Aharonov-Bohm potential $\mathbf{A}_{ab} := \alpha\big(-\frac{x_2}{|x|^2}, \frac{x_1}{|x|^2}\big)$, with $\alpha \in \mathbb{R}$, given by

$$K_{ab}(x,y) = \sum_{k\in\mathbb{Z}} i^{-\beta_k^{ab}} j_{-\alpha_k^{ab}}(|x||y|)\psi_k^{ab}\big(\tfrac{x}{|x|}\big)\overline{\psi_k^{ab}\big(\tfrac{y}{|y|}\big)},$$

where ψ_k^{ab} are the eigenfunctions defined in (2.2) of $L_{\mathbf{A}_{ab},0}$ associated to the eigenvalue $\mu_k^{ab} = (k+\alpha)^2$, and $\alpha_k^{ab}, \beta_k^{ab}$ are given by (1.6) with μ_k replaced by μ_k^{ab}. We have explicitly

$$\alpha_k^{ab} = -\sqrt{\mu_k^{ab}} = -\,|k+\alpha|\,, \qquad \beta_k^{ab} = \sqrt{\mu_k^{ab}} = |k+\alpha|\,.$$

We choose $\alpha = \bar{A}$ with $\bar{A}$ as in (2.9), denote

$$\Sigma^{ab}_{|j|\geqslant\ell}(r,r',\theta,\theta') = \sum_{|j|\geqslant\ell} i^{-|j+\alpha|} j_{|j+\alpha|}(rr')e^{ij\theta}e^{-ij\theta'},$$

and write

$$\Sigma_{|j|\geqslant\ell} = \Big(\Sigma_{|j|\geqslant\ell} - e^{i\bar{A}(\theta-\theta')}\Sigma^{ab}_{|j|\geqslant\ell}\Big) + e^{i\bar{A}(\theta-\theta')}\Sigma^{ab}_{|j|\geqslant\ell}. \tag{3.4}$$

In the paper [16] it has been shown that

$$\sup_{\substack{r,r'\geqslant 0\\ \theta,\theta'\in\mathbb{S}^1}} \Big|e^{i\bar{A}(\theta-\theta')}\Sigma^{ab}_{|j|\geqslant\ell}(r,r',\theta,\theta')\Big| = \sup_{\substack{r,r'\geqslant 0\\ \theta,\theta'\in\mathbb{S}^1}} \Big|\Sigma^{ab}_{|j|\geqslant\ell}(r,r',\theta,\theta')\Big| < \infty. \tag{3.5}$$

To prove the uniform bound of $\Sigma_{|j|\geqslant\ell}$ is hence sufficient to prove the following claim:

$$\sup_{\substack{r,r'\geqslant 0\\ \theta,\theta'\in\mathbb{S}^1}} \left|\Sigma_{|j|\geqslant\ell}(r,r',\theta,\theta') - e^{i\bar{A}(\theta-\theta')}\Sigma^{ab}_{|j|\geqslant\ell}(r,r',\theta,\theta')\right| < \infty. \tag{3.6}$$

In view of the above considerations, we now pass to prove that (3.6) holds.

Let us write

$$\Sigma_{|j|\geqslant\ell} - e^{i\bar{A}(\theta-\theta')}\Sigma^{ab}_{|j|\geqslant\ell} = K_1 + K_2, \tag{3.7}$$

where

$$\begin{aligned} K_1 &= \sum_{|j|\geqslant\ell}\left[i^{-\beta(\lambda_j)}J_{-\alpha(\lambda_j)}(rr') - i^{-|j+\bar{A}|}J_{|j+\bar{A}|}(rr')\right]e^{i(j+\bar{A})\theta}e^{-i(j+\bar{A})\theta'}\\ K_2 &= \sum_{|j|\geqslant\ell} i^{-\beta(\lambda_j)}J_{-\alpha(\lambda_j)}(rr')\times\\ &\qquad\times\left[\left(e^{i(\bar{A}+j)\theta}+R_j(\theta)\right)\left(e^{-i(\bar{A}+j)\theta'}+\overline{R_j(\theta')}\right) - e^{i(j+\bar{A})\theta}e^{-i(j+\bar{A})\theta'}\right]. \end{aligned}$$

Here we used the fact that in dimension $N=2$ we have $j_s \equiv J_s$, for any $s\in\mathbb{R}$.

Let us now recall the estimate

$$|J_\nu(r)| \leqslant \frac{C}{|\nu|^{\frac{1}{3}}} \tag{3.8}$$

(see e.g. [2, 34]), which holds for some $C>0$ independent of x and ν. Moreover, by (2.5) and (2.26) we have that

$$-\alpha(\lambda_j) \sim |j| \qquad \text{as } |j|\to\infty. \tag{3.9}$$

In addition by Lemma 2.1

$$\begin{aligned} \left\|\left(e^{i(\bar{A}+j)\theta}+R_j(\theta)\right)\left(e^{-i(\bar{A}+j)\theta'}+\overline{R_j(\theta')}\right) - e^{i(j+\bar{A})\theta}e^{-i(j+\bar{A})\theta'}\right\|_{L^\infty(\mathbb{S}^1)}&\\ = O\Big(\frac{1}{|j|^3}\Big)& \end{aligned} \tag{3.10}$$

as $|j|\to+\infty$. Hence, by (3.8), (3.9) and (3.10) one easily gets

$$\sup_{\substack{r,r'\geqslant 0\\ \theta,\theta'\in\mathbb{S}^1}} |K_2(r,r',\theta,\theta')| \leqslant C\sum_{|j|\geqslant\ell}|j|^{-\frac{10}{3}} < \infty. \tag{3.11}$$

In order to get the analogous estimate for K_1, we now introduce another well known representation formula for the Bessel functions. Let $\gamma\subset\mathbb{C}$ be the positively oriented contour represented in Figure 1.

Then we have the representation

$$J_\nu(r) = \frac{1}{2\pi i}\int_\gamma e^{\frac{r}{2}\left(z-\frac{1}{z}\right)}\frac{dz}{z^{\nu+1}}$$



FIGURE 1. Integration oriented domain γ.

(see [33, 5.10.7]). Consequently, we obtain

$$
\begin{aligned}
(3.12)\quad & K_1(r,r',\theta,\theta') \\
&= \frac{1}{2\pi i}\sum_{|j|\geqslant \ell}\int_\gamma \frac{1}{z} e^{\frac{rr'}{2}\left(z-\frac{1}{z}\right)}\left[(iz)^{\alpha(\lambda_j)}-(iz)^{-\left|j+\bar{A}\right|}\right] e^{i(j+\bar{A})(\theta-\theta')}\,dz \\
&= \frac{1}{2\pi i}\sum_{|j|\geqslant \ell}\int_\gamma \frac{1}{z} e^{\frac{rr'}{2}\left(z-\frac{1}{z}\right)}(iz)^{-\left|j+\bar{A}\right|}\left[(iz)^{-\sqrt{\lambda_j}+\left|j+\bar{A}\right|}-1\right] e^{i(j+\bar{A})(\theta-\theta')}\,dz.
\end{aligned}
$$

From (2.5) it follows that

$$
(3.13)\quad -\sqrt{\lambda_j}+\left|j+\bar{A}\right| = \sqrt{(j+\bar{A})^2}-\sqrt{\widetilde{a}+(j+\bar{A})^2+O\left(\tfrac{1}{j^2}\right)} = -\frac{\widetilde{a}}{2|j|}+O\left(j^{-2}\right).
$$

Therefore, a first-order Taylor expansion in the last term of (3.12) gives in turn

$$
\begin{aligned}
(3.14)\quad & K_1(r,r',\theta,\theta') \\
&= \frac{1}{2\pi i}\sum_{|j|\geqslant \ell}\int_\gamma \frac{1}{z} e^{\frac{rr'}{2}\left(z-\frac{1}{z}\right)}\left[-\frac{\widetilde{a}\log(iz)}{2|j|}\cdot\frac{e^{i(j+\bar{A})(\theta-\theta')}}{(iz)^{|j+\bar{A}|}}+\mathcal{R}_j(z)\right] dz
\end{aligned}
$$

where $\|\mathcal{R}_j(z)\|_{L^\infty(\gamma)} = O(j^{-2})$ as $|j|\to+\infty$.

We observe that it is possible to exchange the order of summation and integration in (3.14), see the proof of Theorem 1.11 in [16] for details. We hence get

$$
\begin{aligned}
(3.15)\quad & K_1(r,r',\theta,\theta') \\
&= -\frac{1}{2\pi i}\int_\gamma \frac{1}{2z} e^{\frac{rr'}{2}\left(z-\frac{1}{z}\right)}\widetilde{a}\log(iz)\sum_{|j|\geqslant \ell}\left[\frac{1}{|j|}\frac{e^{i(j+\bar{A})(\theta-\theta')}}{(iz)^{|j+\bar{A}|}}+O\left(j^{-2}\right)\right] dz.
\end{aligned}
$$

Finally, we notice that (if ℓ is large enough)

$$\begin{aligned}
&\sum_{|j|\geqslant \ell}^{\infty} \frac{1}{|j|}\frac{e^{i(j+\bar A)(\theta-\theta')}}{(iz)^{|j+\bar A|}}\\
&= -\frac{e^{i\bar A(\theta-\theta')}}{(iz)^{\bar A}}\log\left[1-\frac{e^{i(\theta-\theta')}}{iz}\right] - \frac{e^{i\bar A(\theta-\theta')}}{(iz)^{-\bar A}}\log\left[1-\frac{e^{-i(\theta-\theta')}}{iz}\right]\\
&\quad + \sum_{1\leqslant|j|<\ell}\frac{1}{|j|}\frac{e^{i(j+\bar A)(\theta-\theta')}}{(iz)^{|j|+\bar A\,\mathrm{sgn}\, j}},
\end{aligned}$$

which together with (3.15) leads to

$$\begin{aligned}
&K_1(r,r',\theta,\theta')\\
&= \frac{e^{i\bar A(\theta-\theta')}}{2\pi i}\int_\gamma \frac{1}{2z}e^{\frac{rr'}{2}(z-\frac{1}{z})}\widetilde a\log(iz)\left(\frac{\log\left(1-\frac{e^{i(\theta-\theta')}}{iz}\right)}{(iz)^{\bar A}} + \frac{\log\left(1-\frac{e^{-i(\theta-\theta')}}{iz}\right)}{(iz)^{-\bar A}}\right)\\
&\qquad + \text{bounded terms}.
\end{aligned}$$

In conclusion, since $\left|e^{\frac{r}{2}(z-\frac{1}{z})}\right| = 1$ on Γ_1 and $\log\left(1-\frac{e^{\pm i(\theta-\theta')}}{iz}\right) \sim -\frac{e^{\pm i(\theta-\theta')}}{iz}$ as $|z|\to\infty$, we obtain the desired estimate

$$\sup_{\substack{r,r'\geqslant 0\\ \theta,\theta'\in\mathbb{S}^1}} |K_1(r,r',\theta,\theta')| < \infty, \tag{3.16}$$

which together with (3.7) and (3.11) proves claim (3.6). The proof now follows by (3.3), (3.4), (3.5) and (3.6).

In the resonant case $\widetilde A \in \frac{1}{2}\mathbb{Z}$, we can repeat exactly the same arguments as above, using Lemmas B.9 and B.10 instead of Lemma 2.1; although the control on the remainder terms of the asymptotic expansion is in this case less strong than in the non-resonant case, it is easy to verify that it is enough both for (3.13) and to estimate $\sup|K_2|$ with $C\sum_{|j|\geqslant\ell}|j|^{-\frac{4}{3}} < \infty$ in order to ensure (3.11).

Appendix A. A perturbation lemma

The following result is based on Kato's Perturbation Theory (see [31]).

Lemma A.1. *Let $L_0, L : \mathcal{H}\to\mathcal{H}$ be two self-adjoint operators on a Hilbert space $\mathcal{H}$. Denote by*

$$R_0(\lambda) := (L_0-\lambda I)^{-1} \qquad R(\lambda) := (L-\lambda I)^{-1}.$$

Then:

1. *if $R_0(\lambda), R(\lambda)\in\mathcal{L}(\mathcal{H})$, then λ is not an eigenvalue (neither for R_0 nor for $R(\lambda)$);*
2. *if the operator*

$$T := \frac{1}{2\pi i}\int_\Gamma (R(\lambda)-R_0(\lambda))\,d\lambda$$

has operator norm $\|T\|_{\mathcal{L}(\mathcal{H})} < 1$, being Γ a closed curve in the complex plane, then the number of eigenvalues (counted with multiplicity) of L_0 and L contained in the region bounded by Γ is the same.

Appendix B. Asymptotics of eigenvalues and eigenfunctions in the purely electric case

For the sake of completeness we sketch in this appendix a proof of asymptotics of eigenvalues and eigenfunctions in the purely electric case. We consider the problem

$$\begin{cases} -y'' + a(x)y = \lambda y, & \text{in } [0, 2\pi], \\ y(0) = y(2\pi), \\ y'(0) = y'(2\pi). \end{cases} \tag{B.1}$$

B.1. **The case of a symmetric potential.** Let us first assume that $a(x) \in L^2(\mathbb{S}^1)$ is symmetric with respect to $x = \pi$, i.e.

$$a(\pi - s) = a(\pi + s), \quad \text{for all } s \in [0, \pi]. \tag{B.2}$$

For every $k \in \mathbb{N}$, $k \geqslant 1$, let us consider the space

$$E_k = \{v \in L^2(\mathbb{S}^1) : \ (v, \sin(kt))_{L^2} = 0 \text{ and } (v, \cos(jt))_{L^2} = 0 \text{ for all } j \in \mathbb{N}\}.$$

We denote as $\widetilde{a} = \frac{1}{2\pi}\int_0^{2\pi} a(t)\,dt$ the average of a. For every $j \in \mathbb{N}$ we denote as

$$a_{c,j} = \frac{1}{\pi}\int_0^{2\pi} a(t)\cos(jt)\,dt$$

the j-th cos-Fourier coefficient of $a(x)$.

We observe that since $\int_0^{2\pi}\left(\widetilde{a} - a(x) - \frac{1}{2}a_{c,2k}\right)\sin(kx)\sin(kx)\,dx = 0$ by the definition of $a_{c,2k}$, and $\int_0^{2\pi}\left(\widetilde{a} - a(x) - \frac{1}{2}a_{c,2k}\right)\sin(kx)\cos(kx)\,dx = 0$ by the symmetry condition (B.2), then the problem

$$\begin{cases} -\varphi''(x) - k^2\varphi(x) = \left(\widetilde{a} - a(x) - \frac{1}{2}a_{c,2k}\right)\sin(kx), & \text{in } [0, 2\pi], \\ \varphi(0) = \varphi(2\pi), \\ \varphi'(0) = \varphi'(2\pi), \end{cases} \tag{B.3}$$

admits solutions.

Lemma B.1. *Let $a \in H^{\frac{1}{2}+\delta}([0, 2\pi])$ for some $0 < \delta \ll 1$ such that (B.2) holds. Then, for every $k \in \mathbb{N}$, $k \geqslant 1$, there exists a unique $\varphi_k \in E_k$ solving (B.3). Furthermore there exists some $C > 0$ independent of k such that*

$$\|\varphi_k\|_{L^2} \leqslant C\,\frac{\|a\|_{L^2}}{k}, \quad \|\varphi_k\|_{L^\infty} \leqslant C\,\frac{\|a\|_{H^{\frac{1}{2}+\delta}}}{k}. \tag{B.4}$$

Proof. The unique solution to (B.3) in E_k, i.e. of the form

$$\varphi_k(x) = \sum_{\substack{j=1 \\ j\neq k}}^{\infty} b_j \sin(jx),$$

is given by

$$\varphi_k(x) = \frac{1}{\pi}\sum_{\substack{j=1 \\ j\neq k}}^{\infty} \frac{\int_0^{2\pi}(\widetilde{a} - a(t))\sin(kt)\sin(jt)\,dt}{j^2 - k^2}\sin(jx).$$

We notice that

$$\begin{aligned}\varphi_k(x) &= \frac{1}{\pi}\sum_{\substack{j=1\\ j\neq k}}^{\infty} \frac{\int_0^{2\pi}(\widetilde{a}-a(t))\sin(kt)\sin(jt)\,dt}{j^2-k^2}\sin(jx)\\ &= \frac{1}{2\pi}\sum_{\substack{j=1\\ j\neq k}}^{\infty} \frac{\int_0^{2\pi} a(t)\big(\cos((k-j)t)-\cos((k+j)t)\big)\,dt}{k^2-j^2}\sin(jx)\\ &= \frac{1}{2}\sum_{\substack{j=1\\ j\neq k}}^{\infty} \frac{a_{c,|k-j|}-a_{c,k+j}}{(k-j)(k+j)}\sin(jx).\end{aligned}$$

Therefore

$$\|\varphi_k\|_{L^2}^2 = \frac{\pi}{4}\sum_{\substack{j=1\\ j\neq k}}^{\infty} \frac{\left(a_{c,|k-j|}-a_{c,k+j}\right)^2}{(k-j)^2(k+j)^2} \leqslant \frac{C}{k^2}\,\|a\|_{L^2}^2$$

for some positive constant $C>0$ independent of k. Finally we can estimate

$$\begin{aligned}|\varphi_k(x)| &\leqslant \frac{1}{2}\sum_{\substack{j=1\\ j\neq k}}^{\infty} \frac{|a_{c,|k-j|}|+|a_{c,k+j}|}{|j^2-k^2|}\\ &\leqslant \left(\sum_{j\neq k}|k-j|^{1+2\delta}\left|a_{c,|k-j|}\right|^2\right)^{\frac{1}{2}}\left(\sum_{j\neq k}\frac{1}{|j^2-k^2|^2\,|k-j|^{1+2\delta}}\right)^{\frac{1}{2}}\\ &\quad+\left(\sum_{j\neq k}|k+j|^{1+2\delta}\left|a_{c,|k+j|}\right|^2\right)^{\frac{1}{2}}\left(\sum_{j\neq k}\frac{1}{|j^2-k^2|^2\,|k+j|^{1+2\delta}}\right)^{\frac{1}{2}}\\ &\leqslant \frac{C\|a\|_{H^{\frac{1}{2}+\delta}}}{k}\end{aligned}$$

for some positive constant $C>0$ independent of k. The proof is then complete. □

Lemma B.2. *For every $k\geqslant 1$ and $f\in E_k$ there exists a unique $\tilde{\varphi}_{f,k}\in E_k$ solving*

$$\begin{cases}-\tilde{\varphi}_{f,k}''(x)-k^2\tilde{\varphi}_{f,k}(x)=f(x), & \text{in } [0,2\pi],\\ \tilde{\varphi}_{f,k}(0)=\tilde{\varphi}_{f,k}(2\pi),\\ \tilde{\varphi}_{f,k}'(0)=\tilde{\varphi}_{f,k}'(2\pi).\end{cases}\tag{B.5}$$

Furthermore

$$\|\tilde{\varphi}_{f,k}\|_{L^2}\leqslant\frac{\|f\|_{L^2}}{k},\tag{B.6}$$

$\tilde{\varphi}_{f,k}\in L^\infty(\mathbb{S}^1)$ and for every $\delta\in(0,1)$ there exists $C_\delta>0$ independent of k and f such that

$$\|\tilde{\varphi}_{f,k}\|_{L^\infty}\leqslant\frac{C_\delta}{k^{\frac{1-\delta}{2}}}\,\|f\|_{L^2}\,.\tag{B.7}$$

Proof. Since $f\in E_k$, equation (B.5) is solvable; furthermore, looking for solutions in Fourier series, it turns out that there exists a unique solution in E_k which is

given by

$$\tilde{\varphi}_{f,k}(x) = \sum_{\substack{j=1 \\ j\neq k}}^{\infty} \frac{f_{s,j}\sin(jx)}{j^2-k^2}$$

where

$$f_{s,j} = \frac{1}{\pi}\int_0^{2\pi} f(t)\sin(jt)\,dt.$$

Hence

$$\|\tilde{\varphi}_{f,k}\|_{L^2}^2 = \pi\sum_{\substack{j=1 \\ j\neq k}}^{\infty} \frac{f_{s,j}^2}{(j^2-k^2)^2} \leqslant \frac{\pi}{k^2}\sum_{\substack{j=1 \\ j\neq k}}^{\infty} f_{s,j}^2 = \frac{\|f\|_{L^2}^2}{k^2}$$

and (B.6) is proved. Let $\delta \in (0,1)$. We have that

$$|\tilde{\varphi}_{f,k}(x)| \leqslant \sum_{\substack{j=1 \\ j\neq k}}^{\infty} \frac{|f_{s,j}|}{|j^2-k^2|} \leqslant \Big(\sum_{\substack{j=1 \\ j\neq k}}^{\infty} |f_{s,j}|^2\Big)^{1/2}\Big(\sum_{\substack{j=1 \\ j\neq k}}^{\infty} \frac{1}{|j^2-k^2|^2}\Big)^{1/2} \leqslant C_\delta \frac{\|f\|_{L^2}}{k^{\frac{1-\delta}{2}}}$$

thus completing the proof. □

Proposition B.3. *Let $a \in H^{\frac{1}{2}+\delta}([0,2\pi])$ for some $0 < \delta \ll 1$ such that* (B.2) *holds. Then, for every k sufficiently large, there exists an eigenfunction y_k^s of problem* (B.1) *corresponding to an eigenvalue λ_k^s such that*

$$y_k^s(x) = \sin(kx) + R_k(x), \quad \text{with } \|R_k\|_{L^\infty} = O\Big(\frac{1}{k}\Big) \text{ as } k \to +\infty.$$

$$\lambda_k^s = k^2 + \widetilde{a} - \frac{1}{2}a_{c,2k} + O\Big(\frac{1}{k}\Big) \text{ as } k \to +\infty.$$

Proof. If $k \in \mathbb{N}$, $k \geqslant 1$, and $\varphi \in E_k$, we define

$$\widetilde{\lambda}(\varphi,k) = -\frac{1}{2}a_{c,2k} + \frac{1}{\pi}\int_0^{2\pi} a(x)(\varphi(x)+\varphi_k(x))\sin(kx)dx \tag{B.8}$$

and

$$F(\varphi,k) = \Big(\tfrac{1}{2}a_{c,2k} + \widetilde{\lambda}(\varphi,k)\Big)\sin(kx) + \big(\widetilde{\lambda}(\varphi,k) + \overline{a} - a(x)\big)\big(\varphi(x)+\varphi_k(x)\big). \tag{B.9}$$

Since $\varphi, \varphi_k \in E_k$, from (B.8) it follows that $(F(\varphi,k), \sin(kt))_{L^2} = 0$, while from the symmetry assumption (B.2) we deduce that $(F(\varphi,k), \cos(jt))_{L^2} = 0$ for all j, so that $F(\varphi,k) \in E_k$. Then Lemma B.2 ensures the existence of a unique $T_k(\varphi) \in E_k$ satisfying

$$\begin{cases} -(T_k(\varphi))'' - k^2 T_k(\varphi) = F(\varphi,k), \quad \text{in } [0,2\pi], \\ T_k(\varphi)(0) = T_k(\varphi)(2\pi), \quad T_k(\varphi)'(0) = T_k(\varphi)'(2\pi). \end{cases}$$

Furthermore from (B.6) it follows that

$$\|T_k(\varphi)\|_{L^2} \leqslant \frac{\|F(\varphi,k)\|_{L^2}}{k}.$$

We notice that

$$|\widetilde{\lambda}(\varphi,k)| \leqslant \frac{\|a\|_{L^2}}{\pi}\big(1 + \|\varphi\|_{L^2} + \|\varphi_k\|_{L^2}\big)$$

and, for some $C > 0$ independent of k and φ,

$$\|F(\varphi,k)\|_{L^2} \leqslant C\|a\|_{L^\infty}(1 + \|\varphi\|_{L^2} + \|\varphi_k\|_{L^2})(\|\varphi\|_{L^2} + \|\varphi_k\|_{L^2}). \tag{B.10}$$

The above estimate together with Lemma B.1 imply that, letting

$$B_k = \Big\{\varphi \in E_k : \ \|\varphi\|_{L^2} \leqslant \frac{1}{k^{1-\delta}}\Big\},$$

$T(\varphi) \in B_k$ for every $\varphi \in B_k$. Furthermore it is easy to verify that T is a contraction in B_k endowed with the L^2-metric. Then T admits a unique fixed point in B_k, i.e. there exists a unique $\widetilde{\varphi}_k \in B_k$ such that $T(\widetilde{\varphi}_k) = \widetilde{\varphi}_k$. Then

(B.11)
$$\begin{cases} -\widetilde{\varphi}_k'' - k^2\widetilde{\varphi}_k = \Big(\frac{a_{c,2k}}{2} + \widetilde{\lambda}(\widetilde{\varphi}_k, k)\Big)\sin(kx) + \big(\widetilde{\lambda}(\widetilde{\varphi}_k, k) + \overline{a} - a\big)\big(\widetilde{\varphi}_k + \varphi_k\big), \\ \widetilde{\varphi}_k(0) = \widetilde{\varphi}_k(2\pi), \quad \widetilde{\varphi}_k'(0) = \widetilde{\varphi}_k'(2\pi). \end{cases}$$

From estimate (B.7) of Lemma B.2, (B.10), and Lemma B.1, it follows that, if $\delta \leqslant 1/3$ and k is sufficiently large

$$\begin{aligned} \|\widetilde{\varphi}_k\|_{L^\infty} = \|T(\widetilde{\varphi}_k)\|_{L^\infty} &\leqslant C_\delta \frac{\|F(\widetilde{\varphi}_k, k)\|_{L^2}}{k^{\frac{1-\delta}{2}}} \\ &\leqslant \frac{C_\delta}{k^{\frac{1-\delta}{2}}} C\|a\|_{L^\infty}(1 + \|\widetilde{\varphi}_k\|_{L^2} + \|\varphi_k\|_{L^2})(\|\widetilde{\varphi}_k\|_{L^2} + \|\varphi_k\|_{L^2}) \\ &\leqslant \frac{\text{const}}{k^{\frac{1-\delta}{2}+(1-\delta)}} \leqslant \frac{\text{const}}{k} \end{aligned} \tag{B.12}$$

for some positive const > 0 independent of k. In particular (B.12) implies that

$$\|\widetilde{\varphi}_k\|_{L^2} \leqslant \frac{\text{const}}{k} \tag{B.13}$$

for some positive const > 0 independent of k. From (B.8), (B.13), and (B.4)

$$\widetilde{\lambda}(\widetilde{\varphi}_k, k) + \frac{1}{2}a_{c,2k} = \frac{1}{\pi}\int_0^{2\pi} a(x)(\widetilde{\varphi}_k(x) + \varphi_k(x))\sin(kx)dx = O\Big(\frac{1}{k}\Big) \tag{B.14}$$

as $k \to +\infty$. Let us consider the function

$$y_k(x) = \sin(kx) + \varphi_k(x) + \widetilde{\varphi}_k(x), \quad x \in [0, 2\pi].$$

From the fact that φ_k solves (B.4) and (B.11) we obtain that y_k satisfies

$$\begin{cases} -y_k''(x) + a(x)y_k(x) = (k^2 + \widetilde{a} + \widetilde{\lambda}(\widetilde{\varphi}_k, k))y_k(x), \quad \text{in } [0, 2\pi], \\ y_k(0) = y_k(2\pi), \\ y_k'(0) = y_k'(2\pi), \end{cases} \tag{B.15}$$

i.e. y_k is an eigenfunction of problem (B.1) corresponding to the eigenvalue

$$\lambda_k = k^2 + \widetilde{a} + \widetilde{\lambda}(\widetilde{\varphi}_k, k).$$

From (B.4) and (B.12) we have that $\|y_k(x) - \sin(kx)\|_{L^\infty} = O(\frac{1}{k})$ as $k \to +\infty$, while (B.14) implies that $\lambda_k = k^2 + \widetilde{a} - \frac{1}{2}a_{c,2k} + O(\frac{1}{k})$ as $k \to +\infty$, thus completing the proof. □

Arguing in an analogous way, one can prove the following statement.

Proposition B.4. *Let $a \in H^{\frac{1}{2}+\delta}([0, 2\pi])$ for some $0 < \delta \ll 1$ such that* (B.2) *holds. Then, for every k sufficiently large, there exists an eigenfunction y_k^c of*

problem (B.1) *corresponding to an eigenvalue* λ_k^c *such that*

$$y_k^c(x) = \cos(kx) + R_k(x), \quad \text{with } \|R_k\|_{L^\infty} = O\Big(\frac{1}{k}\Big) \text{ as } k \to +\infty.$$
$$\lambda_k^c = k^2 + \widetilde{a} + \frac{1}{2}a_{c,2k} + O\Big(\frac{1}{k}\Big) \text{ as } k \to +\infty.$$

Remark B.5. If $a \in W^{1,\infty}([0,2\pi])$, then by integrating by parts it follows that

$$|a_{c,2k}| = \frac{1}{2k\pi}\left|\int_0^{2\pi} a'(x)\sin(2kx)dx\right| \leqslant \frac{1}{k}\|a\|_{W^{1,\infty}}.$$

Remark B.6. We notice that, with similar estimates as the ones performed in the proofs of Lemmas B.1 and B.2, the derivative of the remainder terms R_k of Propositions B.3 and B.4 can be estimated as $\|R_k'\|_{L^\infty} = O\big(k^{\frac{1+\delta}{2}}\big)$.

Combining Propositions B.3 and B.4 and Remark B.5, and arguing as in lemmas 2.9 and 2.10 we obtain the following result.

Lemma B.7. *Let* $a \in W^{1,\infty}(\mathbb{S}^1)$ *satisfying* (B.2) *and* $\widetilde{a} = \frac{1}{2\pi}\int_0^{2\pi} a(s)\,ds$. *Let* $\{\mu_k\}_{k\geqslant 1}$ *be the eigenvalues of the operator* $\mathfrak{L}_{0,a}\varphi = -\varphi'' + a\varphi$ *with* 2π*-periodic boundary conditions.*

Then there exist $k^*, \ell \in \mathbb{N}$ *such that* $\{\mu_k : k > k^*\} = \{\lambda_j : j \in \mathbb{Z}, |j| \geqslant \ell\}$ *and*

$$\lambda_j = \widetilde{a} + j^2 + O\big(\tfrac{1}{|j|}\big), \quad \text{as } |j| \to +\infty.$$

Furthermore, for all $j \in \mathbb{Z}$, $|j| \geqslant \ell$, *there exists a* $L^2\big(\mathbb{S}^1,\mathbb{C}\big)$*-normalized eigenfunction* ϕ_j *of the operator* $\mathfrak{L}_{0,a}$ *corresponding to the eigenvalue* λ_j *such that*

$$\phi_j(\theta) = \begin{cases} \frac{1}{\sqrt{\pi}}\Big(\sin(j\theta) + R_j(\theta)\Big), & \text{if } j > 0, \\ \frac{1}{\sqrt{\pi}}\Big(\cos(j\theta) + R_j(\theta)\Big), & \text{if } j < 0, \end{cases}$$

where $\|R_j\|_{L^\infty(\mathbb{S}^1)} = O\big(\frac{1}{|j|}\big)$ *as* $|j| \to \infty$.

B.2. **The general case of a non-symmetric potential.** Let us now drop assumption (B.2). Arguing an in proof of Lemma 2.9, one can prove that there exist some $c > 0$ and $k_0 \in \mathbb{N}$ such that large eigenvalues of $\mathfrak{L}_{0,a}$ are contained in $\bigcup_{j=j_0}^{\infty} B\big(j^2, c\big)$ and, if $j \geqslant j_0$, each ball $B\big(j^2, c\big)$ contains exactly two eigenvalues (counted with their own multiplicity).

Let φ be a $L^2\big(\mathbb{S}^1,\mathbb{C}\big)$-normalized eigenfunction of $\mathfrak{L}_{0,a}$ associated to the eigenvalue $\mu \in B\big(j^2, c\big)$, i.e.

$$\begin{cases} -\varphi''(\theta) + a(\theta)\varphi(\theta) = \mu\varphi(\theta), & \text{in } [0,2\pi], \\ \varphi(0) = \varphi(2\pi), \\ \varphi'(0) = \varphi'(2\pi). \end{cases}$$

From the mean value theorem, there exists $\bar{\theta}_\varphi \in (0,2\pi)$ such that $\varphi'(\bar{\theta}_\varphi) = 0$. The function

$$\widetilde{\varphi}(\theta) = \varphi(\theta + \bar{\theta}_\varphi)$$

satisfies

$$\begin{cases} -\widetilde{\varphi}''(\theta) + a(\theta + \bar{\theta}_\varphi)\widetilde{\varphi}(\theta) = \mu\widetilde{\varphi}(\theta), & \text{in } [0,2\pi], \\ \widetilde{\varphi}(0) = \widetilde{\varphi}(2\pi), \\ \widetilde{\varphi}'(0) = \widetilde{\varphi}'(2\pi) = 0. \end{cases}$$

Hence the function

$$\widehat{\varphi}(\theta)=\begin{cases}\widetilde{\varphi}(2\theta), & \text{if } \theta\in[0,\pi],\\ \widetilde{\varphi}(4\pi-2\theta), & \text{if } \theta\in[\pi,2\pi],\end{cases}$$

is smooth and satisfies

$$\begin{cases}-\widehat{\varphi}''(\theta)+\widehat{a}(\theta)\widehat{\varphi}(\theta)=4\mu\widehat{\varphi}(\theta), & \text{in } [0,2\pi],\\ \widehat{\varphi}(0)=\widehat{\varphi}(2\pi),\\ \widehat{\varphi}'(0)=\widehat{\varphi}'(2\pi)=0,\end{cases}$$

where

$$\widehat{a}(\theta)=\begin{cases}4a(2\theta+\bar{\theta}_\varphi), & \text{if } \theta\in[0,\pi],\\ 4a(4\pi-2\theta+\bar{\theta}_\varphi), & \text{if } \theta\in[\pi,2\pi].\end{cases}$$

We notice that $\widehat{a}\in W^{1,\infty}(\mathbb{S}^1)$, $\widehat{a}$ satisfies (B.2), and $\frac{1}{2\pi}\int_0^{2\pi}\widehat{a}=\frac{4}{2\pi}\int_0^{2\pi}a=4\widetilde{a}$. From Propositions B.3 and B.4 and Lemma B.7, we have that, for some large k,

$$4\mu=k^2+4\widetilde{a}+O(1/|k|) \tag{B.16}$$

and $\widehat{\varphi}$ is a linear combination of the functions $\cos(k\theta)+R_k^1(\theta)$ and $\sin(k\theta)+R_k^2(\theta)$, with $\|R_k^i\|_{L^\infty}=O\big(\frac{1}{k}\big)$ as $k\to+\infty$. The condition $\widehat{\varphi}'(0)=0$ together with Remark B.6 implies that $\widehat{\varphi}$ is a multiple of $\cos(k\theta)+R_k^1(\theta)$, whereas the fact that $\widehat{\varphi}(0)=\widehat{\varphi}(\pi)$ implies that necessarily k is even. From the evenness of k, (B.16), and the fact that $\mu\in B\big(j^2,c\big)$, we conclude that (provided j is large enough) $k=\pm2j$, so that

$$\mu=j^2+\widetilde{a}+O(1/|j|)$$

and $\widehat{\varphi}$ is a multiple of $\cos(2j\theta)+R_{\pm2j}^1(\theta)$. Then

$$\varphi(\theta)=\frac{1}{\sqrt{\pi}}\cos(j(\theta-\bar{\theta}_\varphi))+R_j(\theta)$$

with $\|R_j\|_{L^\infty}=O\big(\frac{1}{|j|}\big)$ as $j\to+\infty$. The above argument proves the following extension to the non-symmetric case of Lemma B.7.

Lemma B.8. *Let $a\in W^{1,\infty}(\mathbb{S}^1)$ and $\widetilde{a}=\frac{1}{2\pi}\int_0^{2\pi}a(s)\,ds$. Let $\{\mu_k\}_{k\geqslant1}$ be the eigenvalues of the operator $\mathfrak{L}_{0,a}\varphi=-\varphi''+a\varphi$ with 2π-periodic boundary conditions. Then there exist $k^*,\ell\in\mathbb{N}$ such that $\{\mu_k:k>k^*\}=\{\lambda_j:j\in\mathbb{Z},|j|\geqslant\ell\}$ and*

$$\lambda_j=\widetilde{a}+j^2+O\big(\tfrac{1}{|j|}\big),\quad \text{as } |j|\to+\infty.$$

Furthermore, for all $j\in\mathbb{Z}$, $|j|\geqslant\ell$, there exist some $\theta_j\in[0,2\pi]$ and a $L^2\big(\mathbb{S}^1,\mathbb{C}\big)$-normalized eigenfunction ϕ_j of the operator $\mathfrak{L}_{0,a}$ on $\mathbb{S}^1$ corresponding to the eigenvalue λ_j such that

$$\phi_j(\theta)=\frac{1}{\sqrt{\pi}}\cos(j(\theta-\theta_j))+R_j(\theta)$$

where $\|R_j\|_{L^\infty(\mathbb{S}^1)}=O\big(\frac{1}{|j|}\big)$ as $|j|\to\infty$.

Combining Lemma B.8 with Lemma 2.3 we obtain the following result which is the analogue of Lemma 2.1 in the case $\widetilde{A}\in\mathbb{Z}$ (we notice that if $\widetilde{A}\in\mathbb{Z}$ then $\bar{A}=0$).

Lemma B.9. *Let $a\in W^{1,\infty}(\mathbb{S}^1)$, $\widetilde{a}=\frac{1}{2\pi}\int_0^{2\pi}a(s)\,ds$, $A\in W^{1,\infty}(\mathbb{S}^1)$ such that $\widetilde{A}=\frac{1}{2\pi}\int_0^{2\pi}A(s)\,ds\in\mathbb{Z}$. Let $\{\mu_k\}_{k\geqslant1}$ be the eigenvalues of the operator $L_{\mathbf{A},a}$ defined in* (1.4) *with* $\mathbf{A}$ *as in* (2.3).

Then there exist $k^*, \ell \in \mathbb{N}$ *such that* $\{\mu_k : k > k^*\} = \{\lambda_j : j \in \mathbb{Z}, |j| \geqslant \ell\}$ *and*

$$\lambda_j = \widetilde{a} + j^2 + O\big(\tfrac{1}{|j|}\big), \quad \text{as } |j| \to +\infty.$$

Furthermore, for all $j \in \mathbb{Z}$, $|j| \geqslant \ell$, *there exist some* $\theta_j \in [0, 2\pi]$ *and a* $L^2\big(\mathbb{S}^1, \mathbb{C}\big)$*-normalized eigenfunction* ϕ_j *of the operator* $L_{\mathbf{A},a}$ *on* $\mathbb{S}^1$ *corresponding to the eigenvalue* λ_j *such that*

$$\phi_j(\theta) = \frac{1}{\sqrt{\pi}} e^{-i\int_0^\theta A(t)\,dt} \cos(j(\theta - \theta_j)) + R_j(\theta),$$

where $\|R_j\|_{L^\infty(\mathbb{S}^1)} = O\big(\frac{1}{|j|}\big)$ *as* $|j| \to \infty$.

B.3. **The case of half-integer circulation.** From the asymptotics of eigenvalues and eigenfunctions in the case of integer circulation discussed above, we can derive the asymptotics of eigenvalues and eigenfunctions also in the case of magnetic potentials with half-integer circulation, corresponding to the case $\widetilde{A} \in \frac{1}{2}\mathbb{Z} \setminus \mathbb{Z}$.

Lemma B.10. *Let* $a \in W^{1,\infty}(\mathbb{S}^1)$, $\widetilde{a} = \frac{1}{2\pi}\int_0^{2\pi} a(s)\,ds$, $A \in W^{1,\infty}(\mathbb{S}^1)$ *such that* $\widetilde{A} = \frac{1}{2\pi}\int_0^{2\pi} A(s)\,ds \in \frac{1}{2}\mathbb{Z} \setminus \mathbb{Z}$. *Let* $\{\mu_k\}_{k\geqslant 1}$ *be the eigenvalues of the operator* $L_{\mathbf{A},a}$ *defined in* (1.4) *with* $\mathbf{A}$ *as in* (2.3). *Then there exist* $k^*, \ell \in \mathbb{N}$ *such that* $\{\mu_k : k > k^*\} = \{\lambda_j : j \in \mathbb{Z}, |j| \geqslant \ell\}$ *and*

$$\lambda_j = \widetilde{a} + \left(j + \frac{1}{2}\right)^2 + O\big(\tfrac{1}{|j|}\big), \quad \text{as } |j| \to +\infty.$$

Furthermore, for all $j \in \mathbb{Z}$, $|j| \geqslant \ell$, *there exist some* $\theta_j \in [0, 2\pi]$ *and a* $L^2\big(\mathbb{S}^1, \mathbb{C}\big)$*-normalized eigenfunction* ϕ_j *of the operator* $L_{\mathbf{A},a}$ *on* $\mathbb{S}^1$ *corresponding to the eigenvalue* λ_j *such that*

$$\phi_j(\theta) = \frac{1}{\sqrt{\pi}} e^{-i\int_0^\theta A(t)\,dt} \Big(\cos\big((j + \tfrac{1}{2})(\theta - \theta_j)\big) + R_j(\theta)\Big),$$

where $\|R_j\|_{L^\infty(\mathbb{S}^1)} = O\big(\frac{1}{|j|}\big)$ *as* $|j| \to \infty$.

Proof. If $\widetilde{A} \in \frac{1}{2}\mathbb{Z} \setminus \mathbb{Z}$, we have that if φ is an eigenfunction of $\mathfrak{L}_{A,a}$ associated to the eigenvalue μ then $\widetilde{\varphi}(\theta) = e^{i\int_0^\theta A(s)\,ds}\varphi(\theta)$ satisfies

$$\begin{cases} -\frac{d^2\widetilde{\varphi}}{d\theta^2} + a(\theta)\widetilde{\varphi} = \mu\widetilde{\varphi} \\ \widetilde{\varphi}(0) = -\widetilde{\varphi}(2\pi) \\ \widetilde{\varphi}'(0) = -\widetilde{\varphi}'(2\pi), \end{cases}$$

and therefore

$$\widehat{\varphi}(\theta) = \begin{cases} \widetilde{\varphi}(2\theta), & \text{if } \theta \in [0, \pi], \\ -\widetilde{\varphi}(2\theta - 2\pi), & \text{if } \theta \in [\pi, 2\pi], \end{cases}$$

satisfies

$$\widehat{\varphi}(s + \pi) = -\widehat{\varphi}(s), \quad \widehat{\varphi}'(s + \pi) = -\widehat{\varphi}'(s) \quad \text{for all } s \in [0, 2\pi] \tag{B.17}$$

and

$$\begin{cases} -\frac{d^2\widehat{\varphi}}{d\theta^2} + \widehat{a}(\theta)\widehat{\varphi} = 4\mu\widehat{\varphi} \\ \widehat{\varphi}(0) = \widehat{\varphi}(2\pi) \\ \widetilde{\varphi}'(0) = \widehat{\varphi}'(2\pi), \end{cases}$$

where

$$\widehat{a}(\theta) = \begin{cases} 4a(2\theta), & \text{if } \theta \in [0,\pi], \\ 4a(2\theta - 2\pi), & \text{if } \theta \in [\pi, 2\pi]. \end{cases}$$

We notice that if $a \in W^{1,\infty}(\mathbb{S}^1)$ then also $\widehat{a} \in W^{1,\infty}(\mathbb{S}^1)$.

From Lemma B.8, we have that, if μ if sufficiently large, then $4\mu = k^2 + 4\widetilde{a} + O(\frac{1}{k})$ for some k large and the eigenspace of the operator $-\frac{d^2}{d\theta^2} + \widehat{a}$ associated to 4μ is generated by one or two functions of the form $\cos(k(\theta - \theta_k)) + R_k(\theta)$ for some $\theta_k \in [0, 2\pi]$ with $\|R_k\|_{L^\infty} = O\big(\frac{1}{k}\big)$ and $\|R_k'\|_{L^\infty} = O\big(k^{\frac{1+\delta}{2}}\big)$. Since condition (B.17) can be satisfied only for odd k, we conclude large eigenvalues of $\mathfrak{L}_{A,a}$ are of the form

$$\left(k + \frac{1}{2}\right)^2 + \widetilde{a} + O\left(\frac{1}{k}\right), \quad \text{as } k \to \infty$$

with eigenfunctions of the form $= e^{-i\int_0^\theta A(s)\,ds} \cos((k + 1/2)(\theta - \theta_k)) + R_k(\theta)$ with $\|R_k\|_{L^\infty} = O\big(\frac{1}{k}\big)$. The conclusion then follows arguing as in lemmas 2.9 and 2.10. □

References

[1] M. Abramowitz and I. A. Stegun, *Handbook of mathematical functions with formulas, graphs, and mathematical tables.* National Bureau of Standards Applied Mathematics Series **55**. For sale by the Superintendent of Documents, U.S. Government Printing Office, Washington, D.C. 1964.

[2] J. A. Barceló, A. Ruiz, and L. Vega, Weighted Estimates for the Helmholtz Equation and Some Applications, *Journal of Functional Analysis* **150** (1997), 356–382.

[3] M. Beceanu and M. Goldberg, Decay estimates for the Schödinger equation with critical potentials, to appear in *Comm. Math. Phys.*, arXiv:1009.5285.

[4] A. Bezubik and A. Strasburger, A new form of the spherical expansion of zonal functions and Fourier transforms of SO(d)-finite functions, *SIGMA Symmetry Integrability Geom. Methods Appl.* **2** (2006), Paper 033, 8 pp.

[5] G. Borg, Umkerhrung der Sturm-Liouvillischen Eigebnvertanfgabe Bestimung der difúerentialgleichung die Eigenverte, *Acta Math.* **78** (1946), 1-96.

[6] Burq, N., Planchon, F., Stalker, J., and Tahvildar-Zadeh, S., Strichartz estimates for the wave and Schrödinger equations with the inverse-square potential, *J. Funct. Anal.* **203** (2003) no. 2, 519–549.

[7] N. Burq, F. Planchon, J. Stalker, and S. Tahvildar-Zadeh, Strichartz estimates for the wave and Schrödinger equations with potentials of critical decay, *Indiana Univ. Math. J.* **53**(6) (2004), 1665–1680.

[8] T. Cazenave, *Semilinear Schrödinger equations.* Courant Lecture Notes in Mathematics **10**, New York University, Courant Institute of Mathematical Sciences, New York; American Mathematical Society, Providence, RI, 2003.

[9] Erdogan, M.B., Goldberg, M., and Schlag, W., Strichartz and Smoothing Estimates for Schrödinger Operators with Almost Critical Magnetic Potentials in Three and Higher Dimensions, *Forum Math.* **21** (2009), 687–722.

[10] M.B. Erdogan, M. Goldberg, and W. Schlag, Strichartz and smoothing estimates for Schrödinger operators with large magnetic potentials in $\mathbb{R}^3$, *J. European Math. Soc.* **10** (2008), 507–531.

[11] P. D'Ancona and L. Fanelli, L^p-boundedness of the wave operator for the one dimensional Schrödinger operators, *Comm. Math. Phys.* **268** (2006), 415–438.

[12] P. D'Ancona and L. Fanelli, Decay estimates for the wave and Dirac equations with a magnetic potential, *Comm. Pure Appl. Math.* **60** (2007), 357–392.

[13] P. D'Ancona and L. Fanelli, Strichartz and smoothing estimates for dispersive equations with magnetic potentials, *Comm. Part. Diff. Eqns.* **33** (2008), 1082–1112.

[14] P. D'Ancona, L. Fanelli, L. Vega, and N. Visciglia, Endpoint Strichartz estimates for the magnetic Schrödinger equation, *J. Funct. Anal.* **258** (2010), 3227–3240.

[15] T. Duyckaerts, Inégalités de résolvante pour l'opérateur de Schrödinger avec potentiel multipolaire critique, *Bulletin de la Société mathématique de France* **134** (2006), 201–239.
[16] L. Fanelli, V. Felli, M. Fontelos, and A. Primo, Time decay of scaling critical electromagnetic Schrödinger flows, *Communications in Mathematical Physics* **324** (2013), 1033–1067.
[17] L. Fanelli and A. García, Counterexamples to Strichartz estimates for the magnetic Schrödinger equation, *Comm. Cont. Math.* **13**(2) (2011), 213–234.
[18] V. Felli, A. Ferrero, and S. Terracini, Asymptotic behavior of solutions to Schrödinger equations near an isolated singularity of the electromagnetic potential, *J. Eur. Math. Soc.* **13** (2011) no. 1, 119–174.
[19] V. Felli, E. M. Marchini, and S. Terracini, On Schrödinger operators with multipolar inverse-square potentials, *Journal of Functional Analysis* **250** (2007), 265–316.
[20] V. Felli, E. M. Marchini, and S. Terracini, On Schrödinger operators with multisingular inverse-square anisotropic potentials, *Indiana Univ. Math. Journal* **58** (2009), 617–676.
[21] V. Georgiev, A. Stefanov, and M. Tarulli, Smoothing - Strichartz estimates for the Schrödinger equation with small magnetic potential, *Discrete Contin. Dyn. Syst.* A **17** (2007), 771–786.
[22] M. Goldberg, Dispersive estimates for the three-dimensional Schrödinger equation with rough potential, *Amer. J. Math.* **128** (2006), 731–750.
[23] M. Goldberg and W. Schlag, Dispersive estimates for Schrödinger operators in dimensions one and three, *Comm. Math. Phys.* **251** (2004) no. 1, 157–178.
[24] M. Goldberg, L. Vega, and N. Visciglia, Counterexamples of Strichartz inequalities for Schrödinger equations with repulsive potentials, *Int. Math Res Not.*, 2006 Vol. 2006: article ID 13927.
[25] J. García Azorero and I. Peral, Hardy inequalities and some critical elliptic and parabolic problems, *J. Differential Equations* **144** (1998), 441–476.
[26] G. Grillo and H. Kovarik, Weighted dispersive estimates for two-dimensional Schrödinger operators with Aharonov-Bohm magnetic field, *Journal of Differential Equations* **256** (2014), 3889–3911.
[27] D. Gurarie, Zonal Schrödinger operators on the n -Sphere: Inverse Spectral Problem and Rigidity, *Comm. Math. Phys.* **131** (1990), 571–603.
[28] G. H. Hardy, J.E. Littlewood, and G. Polya *Inequalities.* Reprint of the 1952 edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 1988.
[29] M. E. H. Ismail, *Classical and quantum orthogonal polynomials in one variable.* Encyclopedia of Mathematics and its Applications, 98. Cambridge University Press, Cambridge, 2005.
[30] H. Kalf, U.-W. Schmincke, J. Walter, R. Wüst, *On the spectral theory of Schrödinger and Dirac operators with strongly singular potentials,* Spectral theory and differential equations (Proc. Sympos., Dundee, 1974; dedicated to Konrad Jörgens), pp. 182–226. Lecture Notes in Math., Vol. 448, Springer, Berlin, 1975.
[31] T. Kato, *Perturbation Theory for Linear Operators*, Springer-Verlag Berlin Heidelberg 1995.
[32] M. Keel and T. Tao, Endpoint Strichartz estimates, *Am. J. Math.* **120** no. 5 (1998), 955–980.
[33] N. N. Lebedev, *Special functions and their applications.* Revised edition, translated from the Russian and edited by Richard A. Silverman. Unabridged and corrected republication. Dover Publications, Inc., New York, 1972.
[34] L. J. Landau, Bessel functions: monotonicity and bounds, *J. London Math. Soc.* **61** (2000), no. 1197–215.
[35] E. H. Lieb and M. Loss, *Analysis,* Graduate Studies in Mathematics 14, AMS (1997).
[36] J. Marzuola, J. Metcalfe, and D. Tataru, Strichartz estimates and local smoothing estimates for asymptotically flat Schrödinger equations, *J. Funct. Anal.* **255** (2008), 1497–1553.
[37] F. Planchon, J. Stalker, and S. Tahvildar-Zadeh, Dispersive estimates for the wave equation with the inverse-square potential, *Discrete Contin. Dyn. Syst.* **9** (2003), 1387–1400.
[38] M. Reed and B. Simon, *Methods of modern mathematical physics. II. Fourier analysis, self-adjointness,* Academic Press, New York-London, 1975.
[39] L. Robbiano and C. Zuily, Strichartz estimates for Schrödinger equations with variable coefficients, Mém. Soc. Math. Fr. (N.S.) **101-102** (2005), vi+208.

[40] I. Rodnianski and W. Schlag, Time decay for solutions of Schrödinger equations with rough and time-dependent potentials, *Invent. Math.* **155** (2004) no. 3, 451–513.
[41] W. Schlag, Dispersive estimates for Schrödinger operators: a survey, *Mathematical aspects of nonlinear dispersive equations, 255285, Ann. of Math. Stud.*, **163**, Princeton Univ. Press, Princeton, NJ, 2007.
[42] B. Simon, Essential self-adjointness of Schrödinger operators with singular potentials, *Arch. Rational Mech. Anal.* **52** (1973), 44–48.
[43] G. Staffilani and D. Tataru, Strichartz estimates for a Schrödinger operator with non-smooth coefficients, *Comm. Partial Differential Equations* **27** (2002) no. 7-8, 1337–1372.
[44] A. Stefanov, Strichartz estimates for the magnetic Schrödinger equation, *Adv. Math.* **210** (2007), 246–303.
[45] L.E. Thomas and C. Villegas-Blas, Singular Continuous Limiting Eigenvalue Distributions for Schrödinger operators on a 2-Sphere, *J. Func. Anal.* **141** (1996), 249–273.
[46] L. E. Thomas and S. R. Wassell, Semiclassical Approximation for Schrödinger operators on a two-sphere at high energy, *J. Math. Phys.* **36** (1995) no. 10, 5480–5505.
[47] R. Weder, The $W_{k,p}$-continuity of the Schrödinger Wave Operators on the line, *Comm. Math. Phys.* **208** (1999), 507–520.
[48] R. Weder, $L^p - L^{p'}$ estimates for the Schrödinger equations on the line and inverse scattering for the nonlinear Schrödinger equation with a potential, *J. Funct. Anal.* **170** (2000), 37–68.
[49] A. Weinstein, Asymptotics for eigenvalue clusters for the laplacian plus a potencial, *Duke Math. J.* **44** (1977), no. 4, 883..892.
[50] K. Yajima, Existence of solutions for Schrödinger evolution equations, *Comm. Math. Phys.* **110** (1987), 415–426.
[51] K. Yajima, The $W^{k,p}$-continuity of wave operators for Schrödinger operators, *J. Math. Soc. Japan* **47** (1995) no. 3, 551–581.
[52] K. Yajima, The $W^{k,p}$-continuity of wave operators for Schrödinger operators III, even dimensional cases $m \geqslant 4$, *J. Math. Sci. Univ. Tokyo* **2** (1995), 311–346.
[53] K. Yajima, L^p-boundedness of wave operators for two-dimensional Schrödinger operators, *Comm. Math. Phys.* **208** (1999) no. 1, 125–152.

Luca Fanelli: SAPIENZA Università di Roma, P.le A. Moro 5, 00185, Roma, Italy
E-mail address: fanelli@mat.uniroma1.it

Veronica Felli: Università di Milano Bicocca, Dipartimento di Matematica e applicazioni, Via Cozzi 55, 20125, Milano, Italy
E-mail address: veronica.felli@unimib.it

Marco Antonio Fontelos: ICMAT-CSIC, Ciudad Universitaria de Cantoblanco. 28049, Madrid, Spain
E-mail address: marco.fontelos@icmat.es

Ana Primo: Departamento de Matemáticas, Universidad Autónoma de Madrid, 28049, Madrid, Spain.
E-mail address: ana.primo@uam.es